

## A GEOMETRIC APPROACH TO PERTURBATION THEORY OF MATRICES AND MATRIX PENCILS. PART I: VERSAL DEFORMATIONS\*

ALAN EDELMAN<sup>†</sup>, ERIK ELMROTH<sup>‡</sup>, AND BO KÄGSTRÖM<sup>‡</sup>

**Abstract.** We derive versal deformations of the Kronecker canonical form by deriving the tangent space and orthogonal bases for the normal space to the orbits of strictly equivalent matrix pencils. These deformations reveal the local perturbation theory of matrix pencils related to the Kronecker canonical form. We also obtain a new singular value bound for the distance to the orbits of less generic pencils. The concepts, results, and their derivations are mainly expressed in the language of numerical linear algebra. We conclude with experiments and applications.

**Key words.** Jordan canonical form, Kronecker canonical form, generalized Schur decomposition, staircase algorithm, versal deformations, tangent and normal spaces, singularity theory, perturbation theory

**AMS subject classifications.** 65F15, 15A21, 15A22

**PII.** S0895479895284634

### Notation.

$\ x\ $	The 2-norm of a vector $x$ .
$A$	A square matrix of size $n \times n$ . $I$ or $I_n$ is the identity matrix.
$A^T$	The transpose of $A$ .
$A^H$	The conjugate transpose of $A$ .
$\bar{A}$	The conjugate of $A$ .
$\ A\ _E$	The Frobenius (or Euclidean) matrix norm.
$\sigma_{\min}(A)$	The smallest singular value of $A$ .
$\text{vec}(A)$	An ordered stack of the columns of a matrix $A$ from left to right.
$\det(A)$	Determinant of $A$ .
$\text{tr}(A)$	Trace of $A$ .
$\ker(A)$	Kernel of space spanned by the columns of $A$ .
$\text{range}(A)$	Range of space spanned by the columns of $A$ .
$\text{diag}(A_1, \dots, A_b)$	A block diagonal matrix with diagonal blocks $A_i$ .
$A \otimes B$	The Kronecker product of two matrices $A$ and $B$ whose $(i, j)$ th block element is $a_{ij}B$ .
$A - \lambda B$	A matrix pencil of size $m \times n$ .
$\lambda_i$	Eigenvalue of $A$ or $A - \lambda B$ . Also, $\gamma_i$ and $\alpha$ are used to denote an eigenvalue.

\*Received by the editors April 17, 1995; accepted for publication (in revised form) by P. Van Dooren July 12, 1996.

<http://www.siam.org/journals/simax/18-3/28463.html>

<sup>†</sup>Department of Mathematics, Room 2-380, Massachusetts Institute of Technology, Cambridge, MA 02139 (edelman@math.mit.edu). The research of this author was supported by NSF grant DMS-9120852 and an Alfred P. Sloan Foundation Research Fellowship.

<sup>‡</sup>Department of Computing Science, Umeå University, S-901 87 Umeå, Sweden (elmroth@cs.umu.se, bokg@cs.umu.se). The research of these authors was supported in part by Swedish National Board of Industrial and Technical Development grant 89-02578P and by Swedish Research Council for Engineering Sciences grant TFR 222-95-34.

$r, r_i, s_i$	$A - \lambda B$ has $r$ distinct eigenvalues $\lambda_i$ of algebraic multiplicity $r_i$ . The sizes of the Jordan blocks associated with an eigenvalue are $s_1 \geq s_2 \geq \dots \geq s_{r_i}$ .
$J_j(\gamma_i)$	Jordan block of size $j \times j$ associated with $\gamma_i$ .
$J_j(\gamma_i, \bar{\gamma}_i)$	Real Jordan block of size $2j \times 2j$ associated with a complex conjugate pair of eigenvalues.
$N_j$	Jordan block of size $j \times j$ associated with the infinite eigenvalue.
$L_j$	Singular block of right (column) minimal index of size $j \times (j + 1)$ .
$L_j^T$	Singular block of left (row) minimal index of size $(j + 1) \times j$ .
$\langle A - \lambda B, C - \lambda D \rangle$	Frobenius inner product of two matrix pairs.
$\text{orbit}(A)$	The set of matrices similar to $A$ .
$\text{orbit}(A - \lambda B)$	The set of matrix pencils equivalent to $A - \lambda B$ .
$\tan(A - \lambda B)$	Tangent space of $\text{orbit}(A - \lambda B)$ at $A - \lambda B$ .
$\text{nor}(A - \lambda B)$	Normal space of $\text{orbit}(A - \lambda B)$ at $A - \lambda B$ .
$\mathcal{S} \oplus \mathcal{T}$	Direct sum of subspaces $\mathcal{S}$ and $\mathcal{T}$ of $\mathbf{R}^n$ .
$\mathcal{S}^\perp$	Subspace perpendicular to $\mathcal{S}$ . $\mathcal{S} \oplus \mathcal{S}^\perp$ is the complete space.
$\dim(\mathcal{S})$	Dimension of subspace $\mathcal{S}$ . $\dim(\mathcal{S})$ denotes dimension of subspace spanned by the columns of $S$ .
$\text{cod}(\mathcal{S})$	Codimension is the dimension of the subspace complementary to $\mathcal{S}$ .
$\mathcal{P}$	The $2mn$ -dimensional space of $m \times n$ matrix pencils, i.e., $\mathcal{P} = \tan(A - \lambda B) \oplus \text{nor}(A - \lambda B)$ .
$\mathcal{V}(p)$	Deformation or (mini)versal deformation with parameter vector $p \in \mathbf{R}^l$ , where $l \geq 1$ . $\mathcal{V}(p)$ is also written $\mathcal{V}(p_1, p_2, \dots, p_l)$ . $q$ is also used as parameter vector.
$Z(p)$	Deformation that spans the orthogonal complement of the orbit of a matrix $A$ .
$Z_A(p) - \lambda Z_B(p)$	Deformation that spans the orthogonal complement of the orbit of a pencil $A - \lambda B$ . Often abbreviated $Z_A - \lambda Z_B$ .

## 1. Introduction and examples.

**1.1. Introduction.** Traditionally, canonical structure computations take as their input some mathematical object, a matrix or a pencil, say, and return an equivalent object that is perhaps simpler or makes clear the structure of the equivalence relation. Some example equivalence relations and corresponding canonical forms are as follows.

Structure	Equivalence relation	Canonical form
Square matrices	$A \sim X^{-1}AX$	Jordan canonical form
Rectangular matrices	$A \sim UAV$	Singular values
Rectangular matrices	$A \sim XA$	Reduced echelon form
Matrix pencils	$A - \lambda B \sim P^{-1}(A - \lambda B)Q$	Kronecker canonical form
Analytic real functions	$f(x) \sim f(\phi(x))$	$\pm x^k$

In the first three examples the input is a matrix. In the next example, the input is a pencil. In these cases,  $X, P$ , and  $Q$  are presumed nonsingular and  $U$  and  $V$  are presumed orthogonal. We presume the real functions  $f$  are analytic in a neighborhood

of zero,  $f(0) = 0$ ,  $\phi(0) = 0$ , and  $\phi(x)$  is monotonic and analytic near zero.

Canonical forms appear in every branch of mathematics. A few examples from control theory may be found in [21, 20, 27, 19]. However, researchers in singularity theory have asked what happens if you have not one object that you want to put into a normal form, but rather a whole family of objects nearby some particular object and you wish to put each member of the family into a canonical form in such a way that the canonical form depends smoothly on the deformation parameters.

For example, one may have a one-parameter matrix deformation of  $A$  which is simply an analytic function  $\mathcal{V}(p)$  for which  $\mathcal{V}(0) = A$ . An  $n$  parameter deformation is defined the same way, except that  $p \in \mathbf{R}^n$ . Similarly, one may have  $n$  parameter deformations of pencils or functions. Remaining with the matrix example, we say two deformations  $\mathcal{V}_1(p)$  and  $\mathcal{V}_2(p)$  are equivalent if  $\mathcal{V}_1(p)$  and  $\mathcal{V}_2(p)$  have the same Jordan canonical form for each and every  $p$ . A deformation of a matrix is said to be versal if, loosely speaking, it captures all possible Jordan form behaviors near the matrix. A deformation is said to be miniversal if it does so with as few parameters as possible. A more formal discussion of these definitions may be found in section 2.

The derivation of versal and miniversal deformations requires a detailed understanding of the perturbation theory of the objects under study. In particular, one needs to understand the tangent space of the equivalence relation and how it is embedded in the entire space. In section 2, we explain the mechanics of this perturbation theory.

While we believe that versal deformations are interesting mathematical objects, this work differs from others on the subject in that our primary goal is not so much the versal deformation or the miniversal deformation, but rather the perturbation theory and how it influences the computation of the Kronecker canonical form. As such, we tend to be interested more in metrical information than topological information. Therefore, we obtain new distance formulas to the space of less generic matrix pencils in section 4. In section 5, we derive an explicit orthogonal basis for the normal space of a Kronecker canonical form. For us a versal decomposition will be an explicit decomposition of a perturbation into its tangential and normal components, and we will not derive any miniversal deformations that may have simpler forms, but hide the metric information.

Versal deformations for function spaces are discussed in [18, 25, 4, 5]. The first application of these ideas for the matrix Jordan canonical form is due to Arnold [1]. Further references closely related to Arnold's matrix approach are [30] and [6]. The latter reference also includes applications to differential equations. Applications of the matrix idea toward an understanding of companion matrix eigenvalue calculations may be found in [13]. The only other work that we are aware of that considers versal deformations of the Kronecker canonical form is by Berg and Kwatny [3], who independently derived some of the normal forms considered in this paper.

Our section 2 contains a thorough explanation of versal deformations from a linear algebra perspective. Section 3 briefly reviews matrix pencils and canonical forms. Section 4 derives the geometry of the tangent and normal spaces to the orbits of matrix pencils. Section 5 derives the versal deformations, while section 6 gives applications and illustrations.

Notation is introduced and defined the first time it appears in the text. Some (but not all) of the notation used in the paper is summarized on the previous page. For example, the glossary of Toeplitz and Hankel matrices (section 5.2) is not repeated there. Moreover, the definitions of different canonical forms (companion, Jordan,

Kronecker, generalized Schur, etc.) are introduced in their context.

**1.2. Geometry of matrix space.** Our guiding message is very simple: matrices should be seen in the mind’s eye geometrically as points in  $n^2$ -dimensional space. A perfect vision of numerical computation would allow us to picture computations as moving matrices from point to point or manifold to manifold.

Abstractly, it hardly matters whether a vector is a column of numbers or a geometric point in space. However, without the interplay of these two representations, numerical linear algebra would not be the same. Imagine explaining without the geometric viewpoint how Householder reflections transform vectors.

In contrast, in numerical linear algebra we all know that matrices are geometric points in  $n^2$ -dimensional space, but it is rare that we actually *think* about them this way. Most often, matrices are thought of as either (sparse or dense) arrays of numbers, or they are operators on vectors.

The Eckart–Young (or Schmidt–Mirsky theorem) [29, p. 210] gives a feel for the geometric approach. The theorem states that the smallest singular value of  $A$  is the Frobenius distance of  $A$  to the set of singular matrices. One can not help but see a blob representing the set of singular matrices. This amorphous blob is most often thought of as an undesirable part of town, so unfortunately numerical analysts hardly ever study the set itself. Algebraic geometers recognize the singular matrices as a variety, meaning that the set can be defined as the zero set of a polynomial system (namely,  $\det(A) = 0$ ). It can also be “stratified” as the union of manifolds. The most generic singular matrices are the ones with rank  $n - 1$ . These matrices form a manifold.

Demmel helped pioneer the development of geometric techniques [7] for the analysis of ill conditioning of numerical analysis problems. Shub and Smale [28] are applying geometrical approaches toward the solution of polynomial systems.

We believe that if only we could better understand the geometry of matrix space, our knowledge of numerical algorithms and their failures would also improve. A general program for numerical linear algebra, then, is to transfer from pure mathematicians the technology to geometrically understand the high dimensional objects that arise in numerical linear algebra. This program may not be easy to follow. A major difficulty is that pure mathematicians pay a price for their beautiful abstractions—they do not always possess a deep understanding of the individual objects that we wish to study. This makes technology transfer difficult. Even when the understanding exists somewhere, it may be difficult to recognize or may be buried under a heavy layer of notation. This makes technology transfer time consuming. Finally, even after expending time excavating, the knowledge may still be difficult to apply toward the understanding or the improving of practical algorithms. This makes technology transfer from pure mathematics frustrating.

Nevertheless, our goal as researchers is the quest for understanding which we may then apply. In this paper, we follow our program for the understanding of the Jordan and Kronecker canonical forms of matrices and matrix pencils, respectively. Many of the ideas in this paper have been borrowed from the pure mathematics literature with the goal of simplifying and applying them to the needs of numerical linear algebraists.

While this is quite a general program for numerical linear algebra, this paper focuses on a particular goal. We analyze *versal deformations* from the numerical linear algebra viewpoint and then compute normal deformations for the Kronecker canonical form. We consider both of these as stepping stones toward the far more difficult goal of truly understanding and improving staircase algorithms for the Jordan

or Kronecker canonical form. These are algorithms used in systems and control theory. The structures of these matrices or pencils reflect important physical properties of the systems they model, such as controllability [10, 32].

The user chooses a parameter  $\eta$  to measure any uncertainty in the data. The existence of a matrix or pencil with a different structure within distance  $\eta$  of the input means that the actual system may have a different structure than the approximation supplied as input. These algorithms try to perturb their input by at most  $\eta$  so as to find a matrix or pencil with as high a codimension as possible. The algorithm is said to *fail* if there is another perturbation of size at most  $\eta$  which would raise the codimension even further. Therefore, we must understand the geometry of matrix space to begin to understand how we can supply the correct information to the user. With this information, we believe that we would then be able to not only correctly provide the least generic solutions, but also understand how singularities hinder this process. Bad solutions may then be refined so as to obtain better solutions. As the next section illustrates, the geometry directly affects the perturbation theory.

**1.3. Motivation: A singular value puzzle.** Consider the following four nearly singular matrices:

$$(1.1) \quad M_1 = \begin{pmatrix} 0 & 1 + \epsilon \\ 0 & 0 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix}, \quad M_3 = \begin{pmatrix} \epsilon & 1 \\ 0 & -\epsilon \end{pmatrix}, \quad M_4 = \begin{pmatrix} \epsilon & 1 \\ 0 & \epsilon \end{pmatrix}.$$

Each of these matrices are distance  $O(\epsilon)$  from the Jordan block

$$J_2(0) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

What is the smaller of the two singular values of each of  $M_1, M_2, M_3$ , and  $M_4$ ? The answer is

$$\sigma_{\min}(M_1) = 0, \quad \sigma_{\min}(M_2) = \epsilon, \quad \sigma_{\min}(M_3) \approx \epsilon^2, \quad \text{and} \quad \sigma_{\min}(M_4) \approx \epsilon^2.$$

A quick way to verify this algebraically is to notice that the larger singular value of each matrix is approximately 1 so that the smaller is approximately the (absolute) determinant of the matrix. Another approach that bounds the smallest singular value is the combination of the Eckart–Young theorem and the observation that these matrices are singular:

$$M'_1 = M_1, \quad M'_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad M'_3 = \begin{pmatrix} \epsilon & 1 \\ -\epsilon^2 & -\epsilon \end{pmatrix}, \quad M'_4 = \begin{pmatrix} \epsilon & 1 \\ \epsilon^2 & \epsilon \end{pmatrix}.$$

When  $\epsilon = 0$  in (1.1) our four matrices become the singular  $2 \times 2$  Jordan block  $J_2(0)$ . As  $\epsilon$  varies from 0 each of the four forms in (1.1) traces out a line in matrix space. The geometric issue that is interesting here is that the line of matrices traced out as  $\epsilon$  varies is {1:In, 2:Normal, 3:Tangent, 4:Tangent} to the set of singular matrices. Somehow, this feels like the “right” explanation for why the smaller singular values are {1:0, 2: $\epsilon$ , 3: $\approx \epsilon^2$ , 4: $\approx \epsilon^2$ }.

Let us take a closer look at the set of singular matrices. The four parameters found in a  $2 \times 2$  matrix  $M$  are best viewed in a transformed coordinate system:

$$M = (x, y, z, w) = x \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + y \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + z \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + w \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} w+z & x \\ y & w-z \end{pmatrix}.$$

In this coordinate system, the singular matrices fall on the surface described by the equation  $w^2 = z^2 + xy$ . This is a three-dimensional surface in four-dimensional space. The traceless singular matrices ( $w = 0$ ) fall on the cone  $z^2 + xy = 0$  in three-dimensional space.

Our matrix  $J_2(0)$  may now be represented as  $(1, 0, 0, 0)$  and the four lines of matrices mentioned above are

$$\begin{aligned} l_1 &= \{(1 + \epsilon, 0, 0, 0)\} = \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \epsilon \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\}, \\ l_2 &= \{(1, \epsilon, 0, 0)\} = \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \epsilon \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right\}, \\ l_3 &= \{(1, 0, \epsilon, 0)\} = \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \epsilon \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right\}, \\ l_4 &= \{(1, 0, 0, \epsilon)\} = \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \epsilon \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\}. \end{aligned}$$

The lines  $l_1, l_2$ , and  $l_3$  are all traceless; i.e., the matrices on each of these lines may be viewed in the three-dimensional space of the cone. The line  $l_1$  is not only tangent to the cone, but in fact it lies in the cone. The line  $l_3$  is tangent to one of the circular cross sections of the cone.




FIG. 1.1. Cone of traceless singular matrices with “stick” representing a tangent.

Figure 1.1 illustrates  $l_3$  as a “stick” resting near the bottom of the cone. The line  $l_1$  is a thin line on the cone through the same point.

The line  $l_4$  is normal to the cone, but it is also tangent to the variety of singular matrices. One way to picture this in three dimensions is to take the three-dimensional slice of  $\{w^2 = z^2 + xy\}$  corresponding to  $x = 1$ , i.e.,  $\{w^2 - z^2 = y\}$ . This is a hyperboloid with the Jordan block as a saddle point. The line is the tangent to the parabola  $w^2 = y$  which rests in the plane  $z = 0$ . Figure 1.2 illustrates this line with a cylindrical stick whose central axis is the tangent. Finally, the line  $l_2$  is normal to the set of singular matrices.

If we move a distance  $\epsilon$  away from a point on a surface along a tangent, our distance to the surface remains  $O(\epsilon^2)$ . This is what the singular value corresponding




FIG. 1.2. Variety of singular matrices. The axis of the cylindrical stick is tangent to the singular variety.

to  $l_3$  and  $l_4$  is telling us. Alternatively, if we move normal to the surface as in  $l_2$ , the singular value changes more rapidly:  $O(\epsilon)$ .

The cone of singular matrices with  $w = 0$  is not only a slice of a large dimensional space, but it is also the (closure of) the set of matrices similar to  $J_2(0)$  (which we denote  $\text{orbit}(J_2(0))$  in section 2.4). The matrices similar to  $J_2(0)$  are singular and traceless. In fact, the only matrix that is singular and traceless that is not similar to  $J_2(0)$  is the 0 matrix which is the vertex of the cone. We further explore this case in section 2.5 after we have defined versal deformations.

We conclude that the geometry of the orbit and, in particular, the directions of the tangents and normals to the orbit directly influence the eigenvalue perturbation theory.

**2. Introduction to versal deformations.** This introduction is designed to be readable for general audiences, but we particularly target the numerical linear algebra community.

The ideas here may be thought of as a numerical analyst's viewpoint on ideas that were inspired by Arnold's work [1] on versal deformations of matrices. Further elaboration upon Arnold's versal deformations of matrices may be found in [6, Chapters 2.9 and 2.10] and [30]. These ideas fit into a larger context of differential topology and singularity theory. Bruce and Giblin [5] have written a wonderfully readable introduction to singularity theory emphasizing the elementary geometrical viewpoint. After reading this introduction, it is easy to be lulled into the belief that one has mastered the subject, but a more advanced wealth of information may be found in [18, 25, 4]. Finally, what none of these references do very well is clearly explain that there is still much in this area that mankind does not yet fully understand.

Singularity theory may be viewed as a branch of the study of curves and surfaces, but its crowning application is toward the topological understanding of functions and their behavior under perturbations. Of course, numerical analysts are very interested in perturbations as well.

### 2.1. Characteristic polynomials give the “feel” of versal deformations.

Let  $\mathcal{V}(p)$  be a differentiable one-parameter family of matrices through  $A \equiv \mathcal{V}(0)$ . This is just a curve in matrix space. If  $A$  has a complicated Jordan canonical form, then very likely the Jordan canonical form of  $\mathcal{V}(p)$  is a discontinuous function of  $p$ . (The Jordan canonical form, you will remember, can have nasty ones popping up unexpectedly on the superdiagonal.) It is even more desirable if that function can somehow describe the kinds of matrices that are near  $A$ .

Discontinuities are as unpleasant for pure mathematicians as they are for computers. Therefore, Arnold [1] asks what kinds of functions of  $p$  are differentiable (or many times differentiable, or analytic).

One function that comes to mind is the characteristic polynomial  $\det(\mathcal{V}(p) - \lambda I)$ . The coefficients of  $\det(\mathcal{V}(p) - \lambda I)$  are clearly differentiable functions of  $p$  no matter how complicated a Jordan canonical form the matrix  $A$  might have. In numerical linear algebra, we never compute the characteristic polynomial because the eigenvalues are often very poorly determined by the coefficients of the characteristic polynomial. Mathematically, the characteristic polynomial is a nice function of a matrix because its coefficients, unlike the eigenvalues of the matrix, are analytic functions of the entries of the matrix.

The characteristic polynomial is a reasonable representation for the Jordan canonical form under the special circumstance that every matrix  $\mathcal{V}(p)$  is *nonderogatory* (i.e., each matrix has exactly one Jordan block for each distinct eigenvalue). By a reasonable representation, we mean here that it actually encodes the Jordan canonical form of  $A$ . Theoretically, if you know the characteristic polynomial, then you know the eigenvalues with appropriate multiplicities. It follows that there is a unique nonderogatory Jordan canonical form (see Wilkinson [35, pp. 11–16 or Note 55, p. 408]). To repeat, there is a one-to-one correspondence among the  $n$  eigenvalues of a nonderogatory matrix, the characteristic polynomial of a nonderogatory matrix, and the Jordan canonical form of a nonderogatory matrix, but only the characteristic polynomial is a differentiable function of the perturbation parameter  $p$ . (The eigenvalues themselves can have first-order perturbations with the nondifferentiable form  $p^{1/n}$ , for example, for an  $n \times n$  matrix  $A$  with only one Jordan block  $J_n(\lambda)$ . This is a well-known example.)

In the language of numerical linear algebra, we would say that a nonderogatory matrix  $A$  may be written in companion matrix form  $KCK^{-1}$  in such a way that differentiable perturbations to the matrix  $A$  lead to differentiable perturbations to the companion matrix  $C$ . Here the matrix  $K$  is a Krylov matrix (see [17, p. 369]). Equivalently, first-order perturbations to the matrix  $A$  are manifested as first-order perturbations to the companion matrix  $C$ . When  $A$  is a companion matrix, this gives a first-order perturbation theory for the characteristic polynomials of nearby matrices. This perturbation theory is computed in [13].

Our story would almost stop here if we were interested only in the Jordan form of nonderogatory matrices. We say “almost” because it would be a shame to stop here without explaining the ideas geometrically. Even if we did not discuss the geometry, we have reasons to continue on, since matrix space is enriched with the derogatory matrices, and also we wish to generalize these ideas about the Jordan canonical form



to cover the more complicated case of the Kronecker canonical form.

### 2.2. The rational canonical form is not enough for derogatory matrices.

In the previous section we saw that  $n$  parameters were sufficient to specify the Jordan canonical form of any matrix in a small neighborhood of a nonderogatory matrix. What happens if the matrix is derogatory? One obvious guess turns out to be wrong. The usual generalization of the companion matrix form for derogatory matrices is the rational canonical form. If  $A$  is derogatory, it may be put in rational canonical form. This form may be thought of as the direct sum of companion matrices  $C_i$  with dimension  $m_1 \geq m_2 \geq \cdots \geq m_k$ . The characteristic polynomial of each  $C_i$  divides the characteristic polynomial of all the preceding  $C_j, j < i$ . Can any nearby matrix be expressed as the direct sum of companion matrices with dimension  $m_1, m_2, \dots, m_k$  in a nice differentiable manner? The answer is generally no; though good enough to specify the Jordan canonical form of a matrix, the rational canonical form fails to be powerful enough to specify the Jordan canonical forms of all matrices in a neighborhood. This is because there are just not enough parameters in the rational canonical form to cover all the possibilities. To have enough parameters we need a “versal deformation.”

One simple example is the identity matrix (or the zero matrix). The rational canonical form has  $m_1 = \cdots = m_n = 1$ . The matrices with this form are the diagonal matrices, and hence every one of them is nondefective (diagonalizable). However, with an arbitrarily small perturbation of the identity, it is possible to obtain defective matrices. The rational canonical form has  $n$  parameters, which are not enough.

**2.3. Versal deformation: The linearized theory.** The “linearized” picture of a versal deformation is easy to understand. We therefore explain this picture before plunging into the global point of view. The general case may be nonlinear, but the linearized theory is all that really matters. For simplicity we assume that we are in real  $n$ -dimensional Euclidean space, but this assumption is not so important.

We recall the elementary fact that if  $\mathcal{S}$  and  $\mathcal{T}$  are subspaces of  $\mathbf{R}^n$  such that  $\mathcal{S} \oplus \mathcal{T} = \mathbf{R}^n$ , then there exist linear projections  $\pi_{\mathcal{S}}$  and  $\pi_{\mathcal{T}}$  that map onto  $\mathcal{S}$  and  $\mathcal{T}$ , respectively.

Consider a point  $x \in \mathcal{S}$ . We will investigate all possible perturbations  $y$  of  $x$ , but we will not be concerned with perturbations that are within  $\mathcal{S}$  itself. Psychologically, we consider all the vectors in  $\mathcal{S}$  to somehow be the same, so there will be no need to distinguish them. Let  $\mathcal{T}$  be any linear subspace such that  $\mathcal{S} \oplus \mathcal{T} = \mathbf{R}^n$ ; i.e., any vector may be written as the sum of an element of  $\mathcal{T}$  and an element of  $\mathcal{S}$  (not necessarily uniquely). Clearly if  $t_1, \dots, t_k$  span  $\mathcal{T}$ , then our perturbed vector  $x + y$  may be written as

$$x + y = x + \sum_{i=1}^k p_i t_i + (\text{something in } \mathcal{S}),$$

where the  $p_i$  may be chosen as linear functions of  $y$ . We see here what will turn out to be the key idea of a versal deformation—every perturbation vector may be expressed in terms of the  $p_i$  and vectors that we are considering to all be equivalent.

We now formally introduce the local picture of versal deformations.

**DEFINITION 2.1.** *A linear deformation of the point  $x$  is a function defined on  $p \in \mathbf{R}^l$ :*

$$\mathcal{V}(p) = x + Tp,$$

where  $T = [t_1, t_2, \dots, t_l]$  are arbitrary directions.

The choice of the word “deformation” is meant to convey the idea that we are looking at small values of the  $p_i$ , and these perturbations are small deformations of the starting point  $x$ .

DEFINITION 2.2. *A linear deformation  $\mathcal{V}_1(p)$  of the point  $x$  is versal if for all linear deformations  $\mathcal{V}_2(q)$  of the point  $x$ , it is possible to write*

$$\mathcal{V}_2(q) = \mathcal{V}_1(\phi(q)) + \theta(q),$$

where  $\phi(q)$  is a linear function from  $q_1, \dots, q_m$  to  $p_1, \dots, p_l$  with  $\phi(0) = 0$  and  $\theta$  is a linear function from  $q$  into  $\mathcal{S}$  with  $\theta(0) = 0$ .

We now explain why  $\mathcal{V}_1(p) = x + \sum_{i=1}^l p_i t_i$  is versal if and only if  $\mathcal{S} \oplus \mathcal{T} = \mathbf{R}^n$ . Clearly  $\mathcal{V}_1(\phi(q)) + \theta(q) \in \mathcal{S} \oplus \mathcal{T}$ , and since  $\mathcal{V}_2(p)$  may be arbitrary, it is necessary that  $\text{span}(\{t_i\}) \oplus \mathcal{S} = \mathbf{R}^n$ . It is also sufficient because we then obtain linear projections allowing us to write  $\mathcal{V}_2(q) = x + \pi_{\mathcal{S}} \mathcal{V}_2(q) + \pi_{\mathcal{T}} \mathcal{V}_2(q)$ . The functions  $\phi$  and  $\theta$  may be obtained from  $\pi_{\mathcal{S}}$  and  $\pi_{\mathcal{T}}$ .

DEFINITION 2.3. *A linear deformation  $\mathcal{V}(p)$  of the point  $x$  is universal or miniversal if it is versal and has the fewest possible parameters needed for a versal deformation.*

The number of parameters in a miniversal deformation is exactly the codimension of  $\mathcal{S}$ . Numerical analysts might prefer taking the  $t_i$  to be an orthogonal basis for  $\mathcal{S}^\perp$ , the subspace perpendicular to  $\mathcal{S}$ . This provides one natural miniversal deformation. Arnold [1] does not insist on using  $\mathcal{S}^\perp$ ; any basis for any subspace of dimension  $n - \dim \mathcal{S}$  will do provided that it intersects  $\mathcal{S}$  at zero only. From the topological point of view, this is exactly the same, though of course the numerical properties may be quite different.

**2.4. Versal deformations—the bigger picture.** The previous section explained the linear or first-order theory of versal deformations. At this point, the reader might wonder whether this is just a whole lot of jargon to merely extend a basis for a subspace to the entire space. At the risk of delaying the motivation until now, we decided to make sure that the linear theory be well understood.

We are still in a finite-dimensional Euclidean space  $\mathbf{R}^n$ , but  $\mathcal{S}$  will no longer be a flat subspace. Instead, we wish to consider any equivalence relation  $\sim$  such that the orbit of  $x$  ( $\text{orbit}(x) \equiv \{y | y \sim x\}$ ) is a sufficiently smooth submanifold. As an example we might define  $x \sim y$  to mean  $\|x\| = \|y\|$ , in which case the orbits are spheres. In this context the word “orbit” is quite natural. In  $n^2$ -dimensional space points may be thought of as  $n \times n$  matrices, and the orbit is the set of matrices with the same Jordan canonical form.

One final example that we must mention (because it explains the origins and significance of singularity theory) lives in an infinite-dimensional space. The vector space is the set of analytic functions  $f(x)$  for which  $f(0) = 0$ . We can define  $f \sim g$  if  $f(x)$  and  $g(\phi(x))$  have the same Taylor expansion at  $x = 0$ , where  $\phi$  is a monotonic analytic function with  $\phi(0) = 0$ . The orbit of any function is some complicated infinite-dimensional manifold, but the codimension of the manifold happens to be finite.

Returning to  $\mathbf{R}^n$ , we can now cast everything into a nonlinear context.

DEFINITION 2.4. *A deformation of the point  $x$  is any continuously differentiable function*

$$\mathcal{V}(p_1, \dots, p_l)$$

satisfying  $\mathcal{V}(0) = x$ .

DEFINITION 2.5. A deformation  $\mathcal{V}_1(p)$  of the point  $x$  is versal if for all deformations  $\mathcal{V}_2(q)$  it is possible to write

$$\mathcal{V}_2(q) \sim \mathcal{V}_1(\phi(q))$$

in an arbitrarily small neighborhood of 0, where  $\phi(q)$  is a continuously differentiable function from  $q_1, \dots, q_m$  to  $p_1, \dots, p_l$  for which  $\phi(0) = 0$ .

The good news is that the inverse function theorem lets us express this nonlinear notion in terms of the linear theory.

THEOREM 2.6. A deformation  $\mathcal{V}(p)$  of  $x$  is versal if and only if  $\mathcal{V}_*(p)$  is a versal linear deformation at the point  $x$  on the subspace  $\tan(\text{orbit}(x))$ , where  $\mathcal{V}_*(p)$  is the linearization of  $\mathcal{V}(p)$  near  $x$  (i.e., only first derivatives matter) and  $\tan$  denotes the subspace tangent to the orbit at  $x$ .

The rigorous proof may be found in [1], but the intuition should be clear: near the point  $x$ , only linear deformations matter, and the curvature of the orbit becomes unimportant—only the tangent plane matters. In other words,  $y \sim x$  only if  $y$  is in the orbit of  $x$ , but to first order  $y \sim x$  if (roughly speaking)  $y = x + s$ , where  $s$  is a small tangent vector to the orbit. The versality theorem (Theorem 2.6) shows that we only have to consider versal linear deformations, which we in the following denote  $\mathcal{V}(p)$ .

**2.5. Versal deformations for the Jordan canonical form.** We begin with deformations of the matrix  $A = J_2(0)$ . The perturbation theory and the normal and tangent spaces were discussed in section 1.3. We will use the same coordinate system here.

Four parameters  $q = (q_1, q_2, q_3, q_4)$  are sufficient to describe the most general deformation of  $A$ :

$$\mathcal{V}_2(q) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} q_1 & q_2 \\ q_3 & q_4 \end{pmatrix}.$$

The equivalence relation is that of similar matrices, and it is easy to see by checking the trace and determinant that for sufficiently small values of  $q$  we have the equivalence

$$\mathcal{V}_2(q) \sim \mathcal{V}_1(p) \equiv \begin{pmatrix} 0 & 1 \\ p_1 & p_2 \end{pmatrix},$$

where  $p = \phi(q)$  is defined by  $p_1 = q_3(1 + q_2) - q_1q_4$  and  $p_2 = q_1 + q_4$ . It is worth emphasizing that the equivalence relation does not work if  $\mathcal{V}_2(q)$  is derogatory, but this does not happen for small parameters  $q$ .

We then see from Definition 2.5 that the two-parameter deformation  $\mathcal{V}_1(p)$  is versal. In fact, it is *miniversal*, in that one needs the two parameters. From the local theory pictured in section 1.3, we saw that the orbit of  $J_2(0)$  is the two-dimensional cone, and therefore the tangent and normal spaces are each two dimensional. The number of parameters in a miniversal deformation is always the dimension of the normal space.

It is a worthwhile exercise to derive the similarity transformation  $S(q)$  (a deformation of the identity matrix) for which

$$\mathcal{V}_2(q) = S(q)^{-1}\mathcal{V}_1(\phi(q))S(q),$$

and then linearize this map for small values of  $q$  to see which directions fall along the tangent space to the cone and which directions are normal to the cone.

Now consider deformations of  $A = I_2$  or  $A = 0$ . Both matrices are derogatory with two eigenvalues 1 and 0, respectively. The tangent space does not exist (i.e., it is zero dimensional). Any possible behavior may be found near  $I_2$  (or 0), including a one-dimensional space of derogatory matrices. The miniversal deformation of  $I_2$  (or 0) is the full deformation requiring four parameters.

The general case has been worked out by Arnold [1]. The tangent vectors to the orbit of a matrix  $A$  are those matrices that may be expressed as  $XA - AX$ . The normal space is the adjoint of the centralizer, i.e., the set of matrices  $Z$  satisfying

$$A^H Z = Z A^H.$$

**DEFINITION 2.7.** *A deformation  $\mathcal{V}(p) = A + Z(p)$  of a matrix  $A$  is a versal deformation if and only if  $Z(p)$  is a basis for the orthogonal complement of  $\text{orbit}(A)$  that intersects the orbit at  $A$ .*

The formal definition of the similarity orbit of a matrix  $A$  is

$$\text{orbit}(A) = \{S^{-1}AS : \det(S) \neq 0\}.$$

The parameterized normal form  $Z(p)$  is the set of matrices that commute with  $A^H$  [1, 16]. For numerical properties we prefer taking  $Z(p)$  to be an orthogonal basis for the normal space of  $\text{orbit}(A)$  at  $A$ . This choice of  $Z(p)$  also ensures that  $\mathcal{V}(p)$  is a miniversal deformation with one parameter for each dimension of the normal space.

Let  $A$  have  $r$  distinct eigenvalues  $\lambda_i, i = 1 : r$  with  $r_i$  Jordan blocks each. Let  $s_1(\lambda_i) \geq s_2(\lambda_i) \geq \dots \geq s_{r_i}(\lambda_i)$  denote the sizes of the Jordan blocks corresponding to the eigenvalue  $\lambda_i$ . Then the dimension of the normal space of  $A$  is

$$\sum_{i=1}^r \sum_{j=1}^{r_i} (2j-1)s_j(\lambda_i) = \sum_{i=1}^r (s_1(\lambda_i) + 3s_2(\lambda_i) + 5s_3(\lambda_i) + \dots).$$

Notice that the values of the distinct  $\lambda_i$  play no role in this formula. The dimension of the normal space of  $A$  is determined only by the sizes of the Jordan blocks of  $A$  associated with distinct eigenvalues. If the matrix is in Jordan canonical form, then the normal space consists of matrices  $Z(p)$  made up of Toeplitz blocks, whose block structure is completely determined by the sizes of the Jordan blocks for different eigenvalues. The normal space is the same for all matrices with the same Jordan structure independent of the values of the distinct eigenvalues, so one may as well consider only Jordan blocks corresponding to a 0 eigenvalue. This form of the normal space for the 0 eigenvalues is a special case in Theorem 5.3.

**3. The algebra of matrix pencils—canonical forms.** We saw in section 2.4 that to consider versal deformations one needs a finite- or infinite-dimensional space and an equivalence relation on this space. For the remainder of this paper, we consider the finite-dimensional Euclidean space of matrix pencils endowed with the Euclidean metric (usually denoted the Frobenius metric in this context). The equivalence relation is that of the strict equivalence of pencils.

We consider a matrix pencil  $A - \lambda B$ , where  $A$  and  $B$  are arbitrary  $m \times n$  matrices with real or complex entries. The pencil is said to be *regular* if  $m = n$  and  $\det(A - \lambda B)$  is not identically zero. Indeed, the zeros of  $\det(A - \lambda B) = 0$  are the (generalized) eigenvalues of a regular pencil. Otherwise, i.e., if  $\det(A - \lambda B)$  is identically zero or

$m \neq n$ ,  $A - \lambda B$  is called *singular*. Two  $m \times n$  pencils  $A_1 - \lambda B_1$  and  $A_2 - \lambda B_2$  are *strictly equivalent* if there exist constant (independent of  $\lambda$ ) invertible matrices  $P$  of size  $m \times m$  and  $Q$  of size  $n \times n$  such that

$$P^{-1}(A_1 - \lambda B_1)Q = A_2 - \lambda B_2.$$

Kronecker has shown that any matrix pencil is strictly equivalent to a canonical diagonal form that describes the structure elements of  $A - \lambda B$  (including generalized eigenvalues and eigenspaces) in full detail (e.g., see [16]). This form is a generalization of the Jordan canonical form (JCF) to general matrix pencils.

**3.1. Kronecker canonical form.** The *Kronecker canonical form* (KCF) of  $A - \lambda B$  exhibits the fine structure elements, including elementary divisors (Jordan blocks) and minimal indices, and is defined as follows [16]. Suppose  $A, B \in \mathbf{C}^{m \times n}$ . Then there exist nonsingular  $P \in \mathbf{C}^{m \times m}$  and  $Q \in \mathbf{C}^{n \times n}$  such that

$$(3.1) \quad P^{-1}(A - \lambda B)Q = \tilde{A} - \lambda \tilde{B},$$

where  $\tilde{A} = \text{diag}(A_1, \dots, A_b)$  and  $\tilde{B} = \text{diag}(B_1, \dots, B_b)$  are block diagonal.  $A_i - \lambda B_i$  is  $m_i \times n_i$ . We can partition the columns of  $P$  and  $Q$  into blocks corresponding to the blocks of  $\tilde{A} - \lambda \tilde{B}$ :  $P = [P_1, \dots, P_b]$ , where  $P_i$  is  $m \times m_i$ , and  $Q = [Q_1, \dots, Q_b]$ , where  $Q_i$  is  $n \times n_i$ . Each block  $M_i \equiv A_i - \lambda B_i$  must be of one of the following forms:  $J_j(\alpha)$ ,  $N_j$ ,  $L_j$ , or  $L_j^T$ . First we consider

$$J_j(\alpha) \equiv \begin{bmatrix} \alpha - \lambda & 1 & & & \\ & \cdot & \cdot & & \\ & & \cdot & \cdot & \\ & & & 1 & \\ & & & & \alpha - \lambda \end{bmatrix} \quad \text{and} \quad N_j \equiv \begin{bmatrix} 1 & -\lambda & & & \\ & \cdot & \cdot & & \\ & & \cdot & \cdot & \\ & & & -\lambda & \\ & & & & 1 \end{bmatrix}.$$

$J_j(\alpha)$  is simply a  $j \times j$  Jordan block, and  $\alpha$  is called a *finite eigenvalue*.  $N_j$  is a  $j \times j$  block corresponding to an *infinite eigenvalue* of multiplicity  $j$ . The  $J_j(\alpha)$  and  $N_j$  blocks together constitute the *regular structure* of the pencil. All the  $A_i - \lambda B_i$  are regular blocks if and only if  $A - \lambda B$  is a regular pencil.  $\sigma(A - \lambda B)$  denotes the eigenvalues of the regular part of  $A - \lambda B$  (with multiplicities) and is called the *spectrum* of  $A - \lambda B$ .

The other two types of diagonal blocks are

$$(3.2) \quad L_j \equiv \begin{bmatrix} -\lambda & 1 & & & \\ & \cdot & \cdot & & \\ & & \cdot & \cdot & \\ & & & -\lambda & 1 \end{bmatrix} \quad \text{and} \quad L_j^T \equiv \begin{bmatrix} -\lambda & & & & \\ 1 & \cdot & & & \\ & \cdot & \cdot & & \\ & & & -\lambda & \\ & & & & 1 \end{bmatrix}.$$

The  $j \times (j+1)$  block  $L_j$  is called a *singular block of right (or column) minimal index  $j$* . It has a one-dimensional right null space  $[1, \lambda, \dots, \lambda^j]^T$  for any  $\lambda$ . The  $(j+1) \times j$  block  $L_j^T$  is a *singular block of left (or row) minimal index  $j$*  and has a one-dimensional left null space for any  $\lambda$ . The left and right singular blocks together constitute the *singular structure* of the pencil and appear in the KCF if and only if the pencil is singular. The regular and singular structures define the *Kronecker structure* of a singular pencil.

We also have a real KCF associated with real matrix pencils. If  $A, B \in \mathbf{R}^{m \times n}$ , there exist nonsingular  $P \in \mathbf{R}^{m \times m}$  and  $Q \in \mathbf{R}^{n \times n}$ , where as before  $P^{-1}(A - \lambda B)Q = \tilde{A} - \lambda \tilde{B}$  is block diagonal. The only difference with (3.1) is the Jordan blocks associated with complex conjugate pairs of eigenvalues. Let  $\alpha = \mu + i\omega$ , where  $\mu, \omega$  are real and

$\omega \neq 0$ . If  $\alpha$  is an eigenvalue of  $A - \lambda B$ , then  $\bar{\alpha}$  is also an eigenvalue. Let  $J_j(\alpha, \bar{\alpha})$  denote a Jordan block of size  $2j \times 2j$  associated with a complex conjugate pair of eigenvalues, here illustrated with the case  $j = 3$ :

$$J_3(\alpha, \bar{\alpha}) \equiv \begin{bmatrix} \mu - \lambda & \omega & 1 & 0 & 0 & 0 \\ -\omega & \mu - \lambda & 0 & 1 & 0 & 0 \\ 0 & 0 & \mu - \lambda & \omega & 1 & 0 \\ 0 & 0 & -\omega & \mu - \lambda & 0 & 1 \\ 0 & 0 & 0 & 0 & \mu - \lambda & \omega \\ 0 & 0 & 0 & 0 & -\omega & \mu - \lambda \end{bmatrix}.$$

The Jordan block  $J_j(\alpha, \bar{\alpha})$  plays the same role in the real JCF as  $\text{diag}(J_j(\alpha), J_j(\bar{\alpha}))$  does in the complex JCF. Notice that each pair of the  $2j$  columns of the real  $P$  and  $Q$  associated with a  $J_j(\alpha, \bar{\alpha})$  block form the real and imaginary parts of the (generalized) principal chains corresponding to the complex conjugate pair of eigenvalues.

**3.2. Generalized Schur form and reducing subspaces.** In most applications it is sufficient to transfer  $A - \lambda B$  to a *generalized Schur form* (e.g., to GUPTRI form [11, 12])

$$(3.3) \quad P^H(A - \lambda B)Q = \begin{bmatrix} A_r - \lambda B_r & * & * \\ 0 & A_{reg} - \lambda B_{reg} & * \\ 0 & 0 & A_l - \lambda B_l \end{bmatrix},$$

where  $P$  ( $m \times m$ ) and  $Q$  ( $n \times n$ ) are unitary and  $*$  denotes arbitrary conforming submatrices. Here the square upper triangular block  $A_{reg} - \lambda B_{reg}$  is regular and has the same regular structure as  $A - \lambda B$  (i.e., contains all eigenvalues (finite and infinite) of  $A - \lambda B$ ). The rectangular blocks  $A_r - \lambda B_r$  and  $A_l - \lambda B_l$  contain the singular structure (right and left minimal indices) of the pencil and are block upper triangular.

$A_r - \lambda B_r$  has only right minimal indices in its KCF, indeed the same  $L_j$  blocks as  $A - \lambda B$ . Similarly,  $A_l - \lambda B_l$  has only left minimal indices in its KCF, the same  $L_j^T$  blocks as  $A - \lambda B$ . If  $A - \lambda B$  is singular at least one of  $A_r - \lambda B_r$  and  $A_l - \lambda B_l$  will be present in (3.3). The explicit structure of the diagonal blocks in staircase form can be found in [12]. If  $A - \lambda B$  is regular  $A_r - \lambda B_r$  and  $A_l - \lambda B_l$  are not present in (3.3) and the GUPTRI form reduces to the upper triangular block  $A_{reg} - \lambda B_{reg}$ . Staircase forms that reveal the Jordan structure of the zero and infinite eigenvalues are contained in  $A_{reg} - \lambda B_{reg}$ .

Given  $A - \lambda B$  in GUPTRI form, we also know different pairs of reducing subspaces [33, 11]. Suppose the eigenvalues on the diagonal of  $A_{reg} - \lambda B_{reg}$  are ordered so that the first  $k$ , say, are in  $\Lambda_1$  (a subset of the spectrum) and the remainder are outside  $\Lambda_1$ . Let  $A_r - \lambda B_r$  be  $m_r \times n_r$ . Then the left and right reducing subspaces corresponding to  $\Lambda_1$  are spanned by the leading  $m_r + k$  columns of  $P$  and leading  $n_r + k$  columns of  $Q$ , respectively. When  $\Lambda_1$  is empty, the corresponding reducing subspaces are called *minimal*, and when  $\Lambda_1$  contains the whole spectrum the reducing subspaces are called *maximal*.

Several authors have proposed (staircase-type) algorithms for computing a generalized Schur form (e.g., see [2, 22, 24, 23, 31, 36]). They are numerically stable in the sense that they compute the exact Kronecker structure (generalized Schur form or something similar) of a nearby pencil  $A' - \lambda B'$ .  $\delta \equiv \|(A - A', B - B')\|_E$  is an upper bound on the distance to the closest  $(A + \delta A, B + \delta B)$  with the KCF of  $(A', B')$ .

Recently, robust software with error bounds for computing the GUPTRI form of a singular  $A - \lambda B$  has been published [11, 12]. Some computational experiments that use this software will be discussed later.

**3.3. Generic and nongeneric Kronecker structures.** Although the KCF looks quite complicated in the general case, most matrix pencils have a quite simple Kronecker structure. If  $A - \lambda B$  is  $m \times n$ , where  $m \neq n$ , then for almost all  $A$  and  $B$  it will have the same KCF, depending only on  $m$  and  $n$ . This corresponds to the *generic case* when  $A - \lambda B$  has full rank for any complex (or real) value of  $\lambda$ . Accordingly, generic rectangular pencils have no regular part. The generic Kronecker structure for  $A - \lambda B$  with  $d = n - m > 0$  is

$$\text{diag}(L_\alpha, \dots, L_\alpha, L_{\alpha+1}, \dots, L_{\alpha+1}),$$

where  $\alpha = \lfloor m/d \rfloor$ , the total number of blocks is  $d$ , and the number of  $L_{\alpha+1}$  blocks is  $m \bmod d$  (which is 0 when  $d$  divides  $m$ ) [31, 8]. The same statement holds for  $d = m - n > 0$  if we replace  $L_\alpha, L_{\alpha+1}$  in (3.2) by  $L_\alpha^T, L_{\alpha+1}^T$ . Square pencils are generically regular; i.e.,  $\det(A - \lambda B) = 0$  if and only if  $\lambda$  is an eigenvalue. The generic singular pencils of size  $n \times n$  have the Kronecker structures [34]

$$\text{diag}(L_j, L_{n-j-1}^T), \quad j = 0, \dots, n - 1.$$

Only if a singular  $A - \lambda B$  is rank deficient (for some  $\lambda$ ) may the associated KCF be more complicated and possibly include a regular part, as well as right and left singular blocks. This situation corresponds to the *nongeneric or degenerate case*, which of course is the real challenge from a computational point of view.

The generic and nongeneric cases can easily be couched in terms of reducing subspaces. For example, generic rectangular pencils have only trivial reducing subspaces and no generalized eigenvalues at all. Generic square singular pencils have the same minimal and maximal reducing subspaces. We think of a nongeneric case as an  $A - \lambda B$  that lies either in a submanifold (its orbit) or the bundle corresponding to similar forms but with differing eigenvalues. In this case the pencil has nontrivial reducing subspaces. Moreover, only if it is perturbed so as to move continuously within this manifold or bundle does its reducing subspaces and generalized eigenvalues also move continuously and satisfy interesting error bounds [9, 11, 14, 26]. These requirements are natural in many control and systems theoretic problems, such as computing controllable subspaces and uncontrollable modes.

**4. The geometry of matrix pencil space.** In the coming sections we derive formulas for the tangent and normal spaces of the orbit of a matrix pencil that we will make use of in computing the versal form in section 5. We also derive new bounds for the distance to less generic pencils.

**4.1. The orbit of a matrix pencil and its tangent and normal spaces.** Any  $m \times n$  matrix pair  $(A, B)$  (with real or complex entries) defines a manifold of *strictly equivalent* matrix pencils in the  $2mn$ -dimensional space  $\mathcal{P}$  of  $m \times n$  pencils:

$$(4.1) \quad \text{orbit}(A - \lambda B) = \{P^{-1}(A - \lambda B)Q : \det(P)\det(Q) \neq 0\}.$$

We may choose a special element of  $\text{orbit}(A - \lambda B)$  that reveals the KCF of the pencil.

As usual, the dimension of  $\text{orbit}(A - \lambda B)$  is equal to the dimension of the tangent space to the orbit at  $A - \lambda B$ , here denoted  $\tan(A - \lambda B)$ . By considering the deformation  $(I_m + \delta X)(A - \lambda B)(I_n - \delta Y)$  of  $A - \lambda B$  to first-order term in  $\delta$ , where  $\delta$  is a

small scalar, we obtain  $A - \lambda B + \delta(X(A - \lambda B) - (A - \lambda B)Y) + O(\delta^2)$ , from which it is evident that  $\tan(A - \lambda B)$  consists of the pencils that can be represented in the form

$$(4.2) \quad T_A - \lambda T_B = (XA - AY) - \lambda(XB - BY),$$

where  $X$  is an  $m \times m$  matrix and  $Y$  is an  $n \times n$  matrix. (This may also be obtained formally by differentiating the exponential map.)

In the language of pure mathematics the map that sends the triple  $(P, Q, A - \lambda B)$  to  $P^{-1}(A - \lambda B)Q$  is called a *group action*. The group is the ordered pair of nonsingular matrices  $(P, Q)$  denoted  $GL_m \times GL_n$  which indicates the size of the matrices and the fact that they are nonsingular. The group  $GL_m \times GL_n$  then is acting on the set of pencils.

A group action is *transitive* if it maps the set onto itself; i.e., if every member of the set may be reached from every other member of the set by the map. Clearly the group action is transitive on orbits. (This is merely a restatement of the definition of an orbit: an orbit is a minimal transitive set with respect to the group action.)

Since the action is transitive, we immediately have that orbits are manifolds. Intuitively, the tangent space “looks” the same at every point, since it may be moved from any point to another point by the group action. Mathematically, the orbit is a *homogeneous space*. The orbit may be equated with the quotient group obtained by forming equivalence classes of pairs  $(P, Q)$  that map  $A - \lambda B$  to the same point. It is a small step to show that reducing subspaces vary smoothly if one perturbs a pencil so that it stays on the same orbit. All one must do is *lift* a curve (maintaining continuity) through a pencil back up to  $GL_m \times GL_n$  and then project out the reducing subspaces.

Using Kronecker products, we can represent the  $2mn$  vectors  $T_A - \lambda T_B \in \tan(A - \lambda B)$  as

$$\begin{bmatrix} \text{vec}(T_A) \\ \text{vec}(T_B) \end{bmatrix} = \begin{bmatrix} A^T \otimes I_m \\ B^T \otimes I_m \end{bmatrix} \text{vec}(X) - \begin{bmatrix} I_n \otimes A \\ I_n \otimes B \end{bmatrix} \text{vec}(Y).$$

In this notation, we may say that the tangent space is the range of the  $2mn \times (m^2 + n^2)$  matrix

$$(4.3) \quad T \equiv \begin{bmatrix} A^T \otimes I_m & -I_n \otimes A \\ B^T \otimes I_m & -I_n \otimes B \end{bmatrix}.$$

We may define the *normal* space  $\text{nor}(A - \lambda B)$  as the space perpendicular to  $\tan(A - \lambda B)$ . Orthogonality in  $\mathcal{P}$ , the  $2mn$ -dimensional space of matrix pencils, is defined with respect to a Frobenius inner product

$$\langle A - \lambda B, C - \lambda D \rangle \equiv \text{tr}(AC^H + BD^H),$$

where  $\text{tr}(X)$  denotes the trace of a square matrix  $X$ . Remembering that the space orthogonal to the range of a matrix is the kernel of the Hermitian transpose, we have that

$$\text{nor}(A - \lambda B) = \ker(T^H) = \ker \begin{bmatrix} \bar{A} \otimes I_m & \bar{B} \otimes I_m \\ -I_n \otimes A^H & -I_n \otimes B^H \end{bmatrix}.$$

In ordinary matrix notation, this states that  $Z_A - \lambda Z_B$  is in the normal space of  $A - \lambda B$  if and only if

$$(4.4) \quad Z_A A^H + Z_B B^H = 0 \quad \text{and} \quad A^H Z_A + B^H Z_B = 0.$$



The conditions on  $Z_A$  and  $Z_B$  can easily be verified and also be derived in terms of the Frobenius inner product, i.e.,

$$(4.5) \quad \langle T_A - \lambda T_B, Z_A - \lambda Z_B \rangle = \text{tr}(X(AZ_A^H + BZ_B^H) - (Z_A^H A + Z_B^H B)Y).$$

*Verification.* If conditions (4.4) are satisfied, it follows from (4.5) that the inner product is zero.

*Derivation.* If  $\langle T_A - \lambda T_B, Z_A - \lambda Z_B \rangle = 0$ , then  $\text{tr}(X(AZ_A^H + BZ_B^H) - (Z_A^H A + Z_B^H B)Y) = 0$  must hold for any  $X$  (of size  $m \times m$ ) and  $Y$  (of size  $n \times n$ ). By choosing  $X \equiv 0$ , (4.5) reduces to  $\text{tr}((Z_A^H A + Z_B^H B)Y) = 0$ , which holds for any  $Y$  if and only if  $Z_A^H A + Z_B^H B = 0$ . Similarly, we can choose  $Y \equiv 0$ , which gives that  $AZ_A^H + BZ_B^H = 0$ .

If  $B = I$ , this reduces to  $Z_A \in \text{nor}(A)$  if and only if  $Z_A^H \in \text{centralizer}(A)$ , which is a well-known fact (e.g., see [1]). We will see in section 5.3 that though the A-part of the normal space is very simple when  $B = I$ , obtaining an orthonormal basis for the B-part is particularly challenging. The requirement that  $Z_B = -A^H Z_A$  when  $B = I$  destroys any orthogonality one may have in a basis for the A-part.

We now collect our general statements and a few obvious consequences.

**THEOREM 4.1.** *Let the  $m \times n$  pencil  $A - \lambda B$  be given. Define the  $2mn \times (m^2 + n^2)$  matrix  $T$  as in (4.3). Then*

$$\tan(A - \lambda B) = \text{range}(T) = \{(XA - AY) - \lambda(XB - BY)\},$$

where  $X$  and  $Y$  are compatible square matrices, and

$$\text{nor}(A - \lambda B) = \ker(T^H) = \{Z_A - \lambda Z_B\},$$

where  $Z_A A^H + Z_B B^H = 0$  and  $A^H Z_A + B^H Z_B = 0$ .

The dimensions of these spaces are

$$(4.6) \quad \dim(\tan(A - \lambda B)) = m^2 + n^2 - \dim(\ker(T))$$

and

$$(4.7) \quad \dim(\text{nor}(A - \lambda B)) = \dim(\ker(T^H)) = \dim(\ker(T)) - (m - n)^2.$$

Of course, the tangent and normal spaces are complementary and span the complete  $2mn$ -dimensional space, i.e.,  $\mathcal{P} = \tan(A - \lambda B) \oplus \text{nor}(A - \lambda B)$ , so that the dimensions in (4.6) and (4.7) add up to  $2mn$ , as they should.

Theorem 4.1 leads to one approach for computing a basis for  $\text{nor}(A - \lambda B)$  from the singular value decomposition (SVD) of  $T$ . Indeed, the left singular vectors corresponding to the zero singular value form such a basis. The dimension of the normal space is also known as the *codimension* of the orbit, here denoted  $\text{cod}(A - \lambda B)$ . Accordingly, we have the following “compact” characterization of the codimension of  $\text{orbit}(A - \lambda B)$ .

**COROLLARY 4.2.** *Let the  $m \times n$  pencil  $A - \lambda B$  be given. Then*

$$\text{cod}(A - \lambda B) = \text{the number of zero singular values of } T.$$

The corresponding result for the (square) matrix case is

$$\text{cod}(A) = \text{the number of zero singular values of } I_n \otimes A - A^T \otimes I_n.$$

Although the SVD-based method is simple and has nice numerical properties (backward stability), it is rather costly in the number of operations. Computing the SVD of  $T$  is an  $O(m^3n^3)$  operation.

Knowing the Kronecker structure of  $A - \lambda B$ , it is also possible to compute the codimension of the orbit as the sum of separate codimensions [8]:

$$(4.8) \quad \text{cod}(A - \lambda B) = c_{\text{Jor}} + c_{\text{Right}} + c_{\text{Left}} + c_{\text{Jor,Sing}} + c_{\text{Sing}}.$$

The different contributions in (4.8) originate from the Jordan structure of all eigenvalues (including any infinite eigenvalue), the right singular blocks ( $L_j \leftrightarrow L_k$ ), the left singular blocks ( $L_j^T \leftrightarrow L_k^T$ ), interactions of the Jordan structure with the singular blocks ( $L_k$  and  $L_j^T$ ), and interactions between the left and right singular structures ( $L_j \leftrightarrow L_k^T$ ), respectively. Explicit expressions for these codimensions are derived in [8]. Assume that the given  $A - \lambda B$  has  $r \leq \min(m, n)$  distinct eigenvalues  $\lambda_i, i = 1 : r$  with  $r_i$  Jordan blocks each. Let  $s_1(\lambda_i) \geq s_2(\lambda_i) \geq \dots \geq s_{r_i}(\lambda_i)$  denote the sizes of the Jordan blocks corresponding to the eigenvalue  $\lambda_i$ . Then the separate codimensions of (4.8) can be expressed as

$$c_{\text{Jor}} = \sum_{i=1}^r \sum_{j=1}^{r_i} (2j-1)s_j(\lambda_i) = \sum_{i=1}^r (s_1(\lambda_i) + 3s_2(\lambda_i) + 5s_3(\lambda_i) + \dots),$$

$$c_{\text{Right}} = \sum_{j>k} (j-k-1), \quad c_{\text{Left}} = \sum_{j>k} (j-k-1), \quad c_{\text{Sing}} = \sum_{j,k} (j+k+2),$$

$$c_{\text{Jor,Sing}} = (\text{size of complete regular part}) \cdot (\text{number of singular blocks}).$$

Notice that if we do not wish to specify the value of an eigenvalue  $\lambda_i$ , the codimension count for this unspecified eigenvalue is one less, i.e.,

$$-1 + s_1(\lambda_i) + 3s_2(\lambda_i) + 5s_3(\lambda_i) + \dots$$

This is sometimes done in algorithms for computing the Kronecker structure of a matrix pencil, where usually only the eigenvalues 0 and  $\infty$  are specified and the remaining ones are unspecified.

It is possible to extract the Kronecker structure of  $A - \lambda B$  from a generalized Schur decomposition in  $O((\max(m, n))^3)$  operations. The most reliable SVD approach for computing a generalized Schur decomposition of  $A - \lambda B$  requires at most  $O((\max(m, n))^4)$  operations, which is still small compared to computing the SVD of  $T$  (4.3) for already moderate values of  $m$  and  $n$  (e.g., when  $m = n$ ).

Speaking loosely, we refer to a pencil as having a particular codimension; when speaking strictly we mean that the orbit of the pencil has this codimension.

For given  $m$  and  $n$  the generic pencil has codimension 0 (i.e., spans the complete  $2mn$ -dimensional space), while the most nongeneric matrix pair  $(A, B) = (0_{m \times n}, 0_{m \times n})$  has codimension =  $2mn$  (i.e., defines a “point” in  $2mn$ -dimensional space). Accordingly, any  $m \times n$  nongeneric pencil different from the “zero pencil” has a codimension  $\geq 1$  and  $< 2mn$ .

**4.2. A lower bound on the distance to a less generic pencil.** The SVD characterization of the codimension of orbit( $A - \lambda B$ ) in Corollary 4.2 leads to the following theorem, from which we present an interesting special case as a corollary.

**THEOREM 4.3.** *For a given  $m \times n$  pencil  $A - \lambda B$  with codimension  $c$ , a lower bound on the distance to the closest pencil  $(A + \delta A) - \lambda(B + \delta B)$  with codimension  $c + d$ , where  $d \geq 1$ , is given by*

$$(4.9) \quad \|(\delta A, \delta B)\|_E \geq \frac{1}{\sqrt{m+n}} \left( \sum_{i=2mn-c-d+1}^{2mn} \sigma_i^2(T) \right)^{1/2},$$

where  $\sigma_i(T)$  denotes the  $i$ th largest singular value of  $T$  ( $\sigma_i(T) \geq \sigma_{i+1}(T) \geq 0$ ).

*Proof.* It follows from Corollary 4.2 that  $T$  has rank =  $2mn - c$  if and only if  $A - \lambda B$  has codimension  $c$  and  $(A + \delta A) - \lambda(B + \delta B)$  has codimension  $c + d$  ( $d \geq 1$ ) if and only if  $T + \delta T$ , where  $\delta T$  is defined as

$$(4.10) \quad \delta T \equiv \begin{bmatrix} \delta A^T \otimes I_m & -I_n \otimes \delta A \\ \delta B^T \otimes I_m & -I_n \otimes \delta B \end{bmatrix},$$

has rank  $2mn - c - d$ . From the construction, it follows that

$$\|\delta T\|_E = \sqrt{m+n} \|(\delta A, \delta B)\|_E$$

(each element  $\delta a_{ij}$  and  $\delta b_{ij}$  appears  $m + n$  times in  $\delta T$ ). The Eckart–Young and Mirsky theorem for finding the closest matrix of a given rank (e.g., see [17]) gives that the size of the smallest perturbation in Frobenius norm that reduces the rank in  $T$  from  $2mn - c$  to  $2mn - c - d$  is

$$(4.11) \quad \left( \sum_{i=2mn-c-d+1}^{2mn-c} \sigma_i^2(T) \right)^{1/2}.$$

Moreover, the fact that  $A - \lambda B$  has codimension  $c$  implies that  $\sigma_{2mn-c+1}(T) = \dots = \sigma_{2mn}(T) = 0$ . Since  $\|\delta T\|_E$  must be larger than or equal to quantity (4.11), the proof is complete.  $\square$

**COROLLARY 4.4.** *For a given generic  $m \times n$  pencil  $A - \lambda B$ , a lower bound on the distance to the closest nongeneric pencil  $(A + \delta A) - \lambda(B + \delta B)$  is given by*

$$(4.12) \quad \|(\delta A, \delta B)\|_E \geq \frac{\sigma_{\min}(T)}{\sqrt{m+n}},$$

where  $\sigma_{\min}(T) = \sigma_{2mn}(T)$  denotes the smallest singular value of  $T$ , which is nonzero for a generic  $A - \lambda B$ .

We remark that the set of  $m \times n$  matrix pencils does not include orbits of all codimensions from 1 to  $2mn$ .

One application of Corollary 4.4 is to characterize the distance to uncontrollability for a multiple input/multiple output linear system  $E\dot{x}(t) = Fx(t) + Gu(t)$ , where  $E$  and  $F$  are  $p \times p$  matrices,  $G$  is  $p \times q$  ( $p \geq q$ ), and  $E$  is assumed to be nonsingular. If  $A - \lambda B \equiv [G|F - \lambda E]$  is generic, the linear system is controllable (i.e., the dimension of the controllable subspace equals  $p$ ) and a lower bound on the distance to the closest uncontrollable system is given by (4.12).

**5. Versal deformations for the KCF.** In the coming sections, we derive versal deformations which for us will mean the decomposition of arbitrary perturbations into the tangent and normal spaces of the orbits of equivalent pencils. Since the set of pencils is itself a vector space, the tangent and normal spaces to the orbits may be thought of as linear affine subplanes embedded in the space of pencils.

DEFINITION 5.1. *A deformation  $\mathcal{V}(p) = A - \lambda B + Z_A(p) - \lambda Z_B(p)$  of a pencil  $A - \lambda B$  is a versal deformation if and only if  $Z_A(p) - \lambda Z_B(p)$  is a basis for the orthogonal complement of  $\text{orbit}(A - \lambda B)$  that intersects the orbit at  $A - \lambda B$ .*

Throughout this paper we will choose  $Z_A(p) - \lambda Z_B(p)$  to have minimum number of parameters and to be an orthogonal basis for the normal space of  $\text{orbit}(A - \lambda B)$  at  $A - \lambda B$ . When it is clear from the context, we will drop the parameters and use the notation  $Z_A - \lambda Z_B$  for the parameterized basis for the normal space.

**5.1. An introductory example.** We start with a small example before considering the general case. Let  $A - \lambda B = L_1 \oplus L_4$  with codimension = 2. (This means that the manifold  $\text{orbit}(A - \lambda B)$  has codimension 2 or dimension 68 in the 70-dimensional space of  $5 \times 7$  pencils.) Since  $A - \lambda B$  is already in KCF we know its block structure:

$$A - \lambda B = \left[ \begin{array}{cc|ccccc} -\lambda & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\lambda & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\lambda & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\lambda & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\lambda & 1 \end{array} \right].$$

From (4.2) the matrices in the tangent space are given by  $T_A - \lambda T_B = (XA - AY) - \lambda(XB - BY)$ , where

$$T_A = \left[ \begin{array}{cc|ccccc} -y_{21} & x_{11} - y_{22} & -y_{23} & x_{12} - y_{24} & x_{13} - y_{25} & x_{14} - y_{26} & x_{15} - y_{27} \\ \boxed{-y_{41}} & x_{21} - y_{42} & -y_{43} & x_{22} - y_{44} & x_{23} - y_{45} & x_{24} - y_{46} & x_{25} - y_{47} \\ -y_{51} & \boxed{x_{31} - y_{52}} & -y_{53} & x_{32} - y_{54} & x_{33} - y_{55} & x_{34} - y_{56} & x_{35} - y_{57} \\ -y_{61} & \boxed{x_{41} - y_{62}} & -y_{63} & x_{42} - y_{64} & x_{43} - y_{65} & x_{44} - y_{66} & x_{45} - y_{67} \\ -y_{71} & x_{51} - y_{72} & -y_{73} & x_{52} - y_{74} & x_{53} - y_{75} & x_{54} - y_{76} & x_{55} - y_{77} \end{array} \right]$$

and

$$T_B = \left[ \begin{array}{cc|ccccc} x_{11} - y_{11} & -y_{12} & x_{12} - y_{13} & x_{13} - y_{14} & x_{14} - y_{15} & x_{15} - y_{16} & -y_{17} \\ x_{21} - y_{31} & -y_{32} & x_{22} - y_{33} & x_{23} - y_{34} & x_{24} - y_{35} & x_{25} - y_{36} & -y_{37} \\ \boxed{x_{31} - y_{41}} & -y_{42} & x_{32} - y_{43} & x_{33} - y_{44} & x_{34} - y_{45} & x_{35} - y_{46} & -y_{47} \\ \boxed{x_{41} - y_{51}} & \boxed{-y_{52}} & x_{42} - y_{53} & x_{43} - y_{54} & x_{44} - y_{55} & x_{45} - y_{56} & -y_{57} \\ x_{51} - y_{61} & \boxed{-y_{62}} & x_{52} - y_{63} & x_{53} - y_{64} & x_{54} - y_{65} & x_{55} - y_{66} & -y_{67} \end{array} \right].$$

By inspection we find the following two relations between elements in  $T_A$  and  $T_B$ :

$$\boxed{\square} : t_{21}^a + t_{32}^a = t_{31}^b + t_{42}^b$$

and

$$\boxed{\square} : t_{31}^a + t_{42}^a = t_{41}^b + t_{52}^b,$$

where  $t_{ij}^a$  and  $t_{ij}^b$  denote the  $(i, j)$ th elements of  $T_A$  and  $T_B$ , respectively. These two relations clearly show that the tangent space has codimension at least 2. It may be verified that the other parameters may be chosen arbitrarily so that the codimension is exactly 2.

We want to find  $Z_A - \lambda Z_B$  that is orthogonal to  $T_A - \lambda T_B$  with respect to the Frobenius inner product, i.e.,

$$(5.1) \quad 0 \equiv \langle T_A - \lambda T_B, Z_A - \lambda Z_B \rangle \equiv \text{tr}(T_A Z_A^H + T_B Z_B^H) \equiv \sum_{i,j} t_{ij}^a \bar{z}_{ij}^a + t_{ij}^b \bar{z}_{ij}^b.$$

This inner product is most easily envisioned as the sum of the elementwise multiplication of the two pencils. Using this point of view, it is obvious that the normal space consists of pencils of the form  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$ :

$$(5.2) \quad Z_A - \lambda Z_B = \left[ \begin{array}{cc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_2 & p_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & p_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right] - \lambda \left[ \begin{array}{cc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -p_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -p_2 & -p_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -p_2 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$= \left[ \begin{array}{cc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_2 + \lambda p_1 & p_1 & 0 & 0 & 0 & 0 & 0 \\ \lambda p_2 & p_2 + \lambda p_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \lambda p_2 & 0 & 0 & 0 & 0 & 0 \end{array} \right],$$

where  $p_1$  and  $p_2$  are arbitrary. Roughly speaking, the parameter  $p_1$  corresponds to the doubly boxed entries ( $\boxed{\square}$ ) and the parameter  $p_2$  corresponds to the singly boxed entries. ( $\square$ ).

Now,  $\mathcal{V}(p) = A - \lambda B + Z_A - \lambda Z_B$  may be thought of as a versal deformation, or normal form, with minimum number of parameters (equal to the codimension of the original pencil). It follows that any (complex) pencil close to the given  $A - \lambda B$  in KCF can be reduced to the two-parameter normal form  $\mathcal{V}(p) = A - \lambda B + Z_A - \lambda Z_B$  in terms of equivalence transformations that are deformations of the identity.

**5.2. Notation: A glossary of Toeplitz and Hankel matrices.** The example in the previous section shows that a nonzero block of  $Z_A - \lambda Z_B$  has a structured form. Indeed, the  $(2, 1)$  block has a Toeplitz-like form with  $j - i = 3$  nonzero diagonals starting from the  $(1, 1)$  element of the  $(2, 1)$  block. A closer look shows that the A-part has  $i - j - 1 = 2$  nonzero diagonals and the B-part is just the same matrix negated and with the diagonals shifted one row downward. In general, different nonzero blocks with Toeplitz or Hankel properties will show up in  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$ . To simplify the proof of the general case we introduce some Toeplitz and Hankel matrices. Arrows and “stops” near the matrices make clear how the matrix is defined.

Let  $S_{s \times t}^L$  be a lower trapezoidal  $s \times t$  Toeplitz matrix with the first nonzero diagonal

starting at position  $(1, 1)$ :

$$S_{s \times t}^L = \begin{matrix} \downarrow \\ \left[ \begin{array}{ccc} p_1 & 0 & 0 \\ \vdots & \ddots & 0 \\ \vdots & & p_1 \\ p_{s-t+1} & & \vdots \\ \vdots & \ddots & \vdots \\ \perp \left[ \begin{array}{ccc} p_s & \cdots & p_{s-t+1} \end{array} \right] \end{array} \right. \end{matrix} \text{ if } s \geq t \text{ and } S_{s \times t}^L = \begin{matrix} \downarrow \\ \left[ \begin{array}{cccc} p_1 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ \perp \left[ \begin{array}{cccc} p_s & \cdots & p_1 & 0 \cdots 0 \end{array} \right] \end{array} \right. \end{matrix} \text{ otherwise,}$$

and let  $T_{s \times t}^L$  be a *lower trapezoidal*  $s \times t$  Toeplitz matrix with the first nonzero diagonal's last element at position  $(s, t)$ :

$$T_{s \times t}^L = \begin{matrix} \left[ \begin{array}{ccc} 0 & \cdots & 0 \\ \vdots & & \vdots \\ 0 & & \vdots \\ p_1 & \ddots & \vdots \\ \vdots & \ddots & 0 \\ \perp \left[ \begin{array}{ccc} p_t & \cdots & p_1 \end{array} \right] \end{array} \right. \end{matrix} \text{ if } s \geq t \text{ and } T_{s \times t}^L = \begin{matrix} \left[ \begin{array}{cccc} p_{t-s+1} & \cdots & p_1 & 0 & 0 \\ \vdots & \ddots & & \ddots & 0 \\ \perp \left[ \begin{array}{cccc} p_t & \cdots & p_{t-s+1} & \cdots & \leftarrow p_1 \end{array} \right] \end{array} \right. \end{matrix} \text{ otherwise.}$$

If  $s < t$ , the entries of the last  $t - s$  columns of  $S_{s \times t}^L$  are zero. Similarly, if  $s \geq t$ , the entries of the first  $s - t$  rows of  $T_{s \times t}^L$  are zero.

Let  $S_{s \times t}^B$  be a *banded lower trapezoidal*  $s \times t$  Toeplitz with last row 0:

$$S_{s \times t}^B = \begin{matrix} \downarrow \\ \left[ \begin{array}{ccc} p_1 & 0 & 0 \\ \vdots & \ddots & 0 \\ \vdots & & p_1 \\ \perp \left[ \begin{array}{ccc} p_{s-t} & & \vdots \\ 0 & \ddots & \vdots \\ \vdots & \ddots & p_{s-t} \\ 0 & \cdots & 0 \end{array} \right] \end{array} \right. \end{matrix} \text{ if } s > t \text{ and } S_{s \times t}^B = 0 \text{ otherwise,}$$

and let  $T_{s \times t}^B$  be another *banded lower trapezoidal*  $s \times t$  Toeplitz matrix, this time with last column 0:

$$T_{s \times t}^B = \begin{matrix} \left[ \begin{array}{cccc} \leftarrow p_{t-s} & \cdots & \leftarrow p_1 & 0 \cdots 0 \\ 0 & \ddots & & \ddots & \vdots \\ 0 & 0 & p_{t-s} & \cdots & p_1 & 0 \end{array} \right. \end{matrix} \text{ if } s < t \text{ and } T_{s \times t}^B = 0 \text{ otherwise.}$$

Notice that the last row of  $S_{s \times t}^B$  (if  $s > t$ ) and the last column of  $T_{s \times t}^B$  (if  $s < t$ ) have all entries equal to zero.

Moreover, let  $H_{s \times t}^L$  be a *lower trapezoidal*  $s \times t$  Hankel matrix with the first nonzero diagonal starting at position  $(1, t)$ :

$$H_{s \times t}^L = \begin{bmatrix} 0 & 0 & p_1 & & \\ 0 & \ddots & \vdots & & \\ p_1 & & \vdots & & \\ \vdots & & p_{s-t+1} & & \\ \vdots & \ddots & \vdots & & \\ p_{s-t+1} & \cdots & p_s & & \end{bmatrix} \begin{matrix} \downarrow \\ \\ \\ \\ \perp \end{matrix} \quad \text{if } s \geq t \text{ and } H_{s \times t}^L = \begin{bmatrix} 0 & \cdots & 0 & p_1 \\ \vdots & & \ddots & \vdots \\ 0 \cdots 0 & p_1 & \cdots & p_s \end{bmatrix} \begin{matrix} \downarrow \\ \\ \perp \end{matrix} \text{ otherwise,}$$

and let  $H_{s \times t}^U$  be a similar *upper trapezoidal*  $s \times t$  Hankel matrix:

$$H_{s \times t}^U = \begin{bmatrix} \vdash & \leftarrow & & & \\ p_t & \cdots & p_1 & & \\ \vdots & \ddots & 0 & & \\ p_1 & \ddots & \vdots & & \\ 0 & & \vdots & & \\ \vdots & & \vdots & & \\ 0 & \cdots & 0 & & \end{bmatrix} \begin{matrix} \\ \\ \\ \\ \\ \end{matrix} \quad \text{if } s \geq t \text{ and } H_{s \times t}^U = \begin{bmatrix} \vdash & & & \leftarrow & \\ p_t & \cdots & p_{t-s+1} & \cdots & p_1 \\ \vdots & \ddots & & \ddots & 0 \\ p_{t-s+1} & \cdots & p_1 & 0 & 0 \end{bmatrix} \begin{matrix} \\ \\ \\ \end{matrix} \text{ otherwise.}$$

If  $s < t$ , the entries of the first  $t - s$  columns of  $H_{s \times t}^L$  are zero. Similarly, if  $s \geq t$ , the entries of the last  $s - t$  rows of  $H_{s \times t}^U$  are zero.

Let  $H_{s \times t}$  be a *dense*  $s \times t$  Hankel matrix (with the first diagonal starting at position  $(1, 1)$ ):

$$H_{s \times t} = \begin{bmatrix} p_1 & p_2 & p_3 & \cdots & p_t \\ p_2 & \ddots & & & \vdots \\ p_3 & & & & \vdots \\ \vdots & & & & \vdots \\ p_s & \cdots & p_{s+t-1} & & \end{bmatrix}$$

for both the cases  $s \geq t$  and  $s < t$ .

The *nilpotent*  $k \times k$  matrix

$$C_k = \begin{bmatrix} 0 & I_{k-1} \\ 0 & 0 \end{bmatrix}$$

will be used as a shift operator. For a given  $k \times n$  matrix  $X$ , the rows are shifted one row upward and downward by the operations  $C_k X$  and  $C_k^T X$ , respectively. The columns are shifted one column rightward and one column leftward in an  $n \times k$  matrix  $X$  by the operations  $X C_k$  and  $X C_k^T$ , respectively. The  $k \times (k + 1)$  matrices

$$G_k = [I_k \ 0] \text{ and } \hat{G}_k = [0 \ I_k],$$

will be used to pick all rows but one or all columns but one of a given matrix  $X$  in the following way. The first  $k$  and last  $k$  rows in a  $(k + 1) \times n$  matrix  $X$  are picked by

$G_k X$  and  $\hat{G}_k X$ , respectively. The  $k$  first and  $k$  last columns in an  $n \times (k + 1)$  matrix  $X$  are picked by  $XG_k^T$  and  $X\hat{G}_k^T$ , respectively.

Let  $\hat{I}_k$  denote the  $k \times k$  matrix obtained by reversing the order of the columns in the  $k \times k$  identity matrix. It follows that for an  $n \times k$  matrix  $X$ , the order of the columns is reversed by the multiplication  $X\hat{I}_k$ .

So far, the matrices introduced are rectangular Toeplitz and Hankel matrices with a special structure, e.g., lower trapezoidal ( $S^L, T^L, H^L$ ), banded lower trapezoidal ( $S^B, T^B$ ), upper trapezoidal ( $H^U$ ), or dense ( $H$ ). The matrices  $C$  and  $G, \hat{G}$  that will be used as “shift” and “pick” operators, respectively, are Toeplitz matrices with only one nonzero diagonal. In the next section we will see that versal deformations for all combinations of different blocks in the KCF, except Jordan blocks with nonzero finite eigenvalues, can be expressed in terms of these matrices. To cope with nonzero finite Jordan blocks  $J_k(\gamma), \gamma \neq 0$  we need to introduce three more matrices. First, we introduce two lower triangular Toeplitz matrices  $D^L$  and  $E^L$ , which are involved in the case with two  $J_k(\gamma)$  blocks. Finally, we introduce the “monstrous” matrix  $F^D$ , which captures the cases with a (left or right) singular block and a  $J_k(\gamma)$  block.

Given  $\gamma \neq \{0, \infty\}$ , define two infinite sequences of numbers  $d_i$  and  $e_i$  by the recursion

$$(5.3) \quad \begin{bmatrix} d_i \\ \gamma e_i \end{bmatrix} = - \begin{bmatrix} 1 & 1 \\ 1 & 2 - 1/i \end{bmatrix} \begin{bmatrix} \bar{\gamma} d_{i-1} \\ e_{i-1} \end{bmatrix}$$

starting with

$$\begin{bmatrix} d_1 \\ e_1 \end{bmatrix} = \begin{bmatrix} \gamma \\ 1 \end{bmatrix}.$$

Given sizes  $s$  and  $t$ , for  $1 \leq q \leq \min\{s, t\}$ , we define  $D_{s \times t}[q]$  and  $E_{s \times t}[q]$  as lower triangular Toeplitz matrices with  $q$  diagonals in terms of  $d_1, \dots, d_q$  and  $e_1, \dots, e_{q-1}$  and a boundary value  $e_q^* = -\bar{\gamma}d_q$ :

$$D_{s \times t}[q] = \begin{bmatrix} 0 & \dots & 0 \\ d_q & & \\ d_{q-1} & \ddots & \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ d_2 & & \ddots & \ddots & \\ d_1 & d_2 & \dots & d_{q-1} & d_q & 0 \end{bmatrix} \quad \text{and} \quad E_{s \times t}[q] = \begin{bmatrix} 0 & \dots & 0 \\ e_q^* & & \\ e_{q-1} & \ddots & \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ e_2 & & \ddots & \ddots & \\ e_1 & e_2 & \dots & e_{q-1} & e_q^* & 0 \end{bmatrix}.$$

We take linear combinations with parameters  $p_j$  to form the matrices

$$(5.4) \quad D_{s \times t}^L = \sum_{i=1}^{\min\{s,t\}} p_j D_{s \times t}[i] \pi(i) \quad \text{and} \quad E_{s \times t}^L = \sum_{i=1}^{\min\{s,t\}} p_j E_{s \times t}[i] \pi(i),$$

where  $j = \min\{s, t\} - i + 1$  and  $\pi(i) = -\prod_{k=2}^{i-1} k\gamma/(1-2k)$  is defined to be  $1/\gamma$  and  $-1$  for  $i = 1$  and  $i = 2$ , respectively. The parameter index  $j$  and the scaling function  $\pi(i)$  are chosen to satisfy  $D_{s \times t}^L = S_{s \times t}^L$  and  $E_{s \times t}^L = -C_s^T S_{s \times t}^L$  for  $\gamma = 0$  in Theorem 5.3 (see Tables 5.1 and 5.2). By simplifying (5.4) using  $i = j$  and  $\pi(i) = 1$ , this consistency will be lost, but we will still have valid expressions for the versal deformations.



The relations between the elements of  $D_{s \times t}^L$  and  $E_{s \times t}^L$  are most readily shown by an example:

$$D_{4,3}^L = \begin{bmatrix} 0 & 0 & 0 \\ p_1 \left( \frac{2|\gamma|^4}{3} + \frac{4|\gamma|^2}{3} + 1 \right) & 0 & 0 \\ p_1 \left( -\frac{2\bar{\gamma}|\gamma|^2}{3} - \frac{2\bar{\gamma}}{3} \right) + p_2 (|\gamma|^2 + 1) & p_1 \left( \frac{2|\gamma|^4}{3} + \frac{4|\gamma|^2}{3} + 1 \right) & 0 \\ p_1 \frac{2\bar{\gamma}^2}{3} - p_2 \bar{\gamma} + p_3 & p_1 \left( -\frac{2\bar{\gamma}|\gamma|^2}{3} - \frac{2\bar{\gamma}}{3} \right) + p_2 (|\gamma|^2 + 1) & p_1 \left( \frac{2|\gamma|^4}{3} + \frac{4|\gamma|^2}{3} + 1 \right) \end{bmatrix}$$

and

$$E_{4,3}^L = \begin{bmatrix} 0 & 0 & 0 \\ p_1 \left( -\frac{2\bar{\gamma}|\gamma|^4}{3} - \frac{4\bar{\gamma}|\gamma|^2}{3} - \bar{\gamma} \right) & 0 & 0 \\ p_1 \left( -\frac{2|\gamma|^2}{3} - 1 \right) + p_2 (-\bar{\gamma}|\gamma|^2 - \bar{\gamma}) & p_1 \left( -\frac{2\bar{\gamma}|\gamma|^4}{3} - \frac{4\bar{\gamma}|\gamma|^2}{3} - \bar{\gamma} \right) & 0 \\ p_1 \frac{2\bar{\gamma}}{3} - p_2 - p_3 \bar{\gamma} & p_1 \left( -\frac{2|\gamma|^2}{3} - 1 \right) + p_2 (-\bar{\gamma}|\gamma|^2 - \bar{\gamma}) & p_1 \left( -\frac{2\bar{\gamma}|\gamma|^4}{3} - \frac{4\bar{\gamma}|\gamma|^2}{3} - \bar{\gamma} \right) \end{bmatrix}.$$

Let  $F_{s \times t}^D$  ( $D$  for dense) be defined as

$$F_{s \times t}^D = \sum_{i=1}^s p_{s-i+1} F_{s \times t}[i],$$

where  $F_{s \times t}[q]$  has the  $q$  last rows nonzero and defined as

$$(5.5) \quad \begin{aligned} f_{s-q+1,j} &= \bar{\gamma}^{j-1} && \text{for } j = 1, \dots, t, \\ f_{i,j} &= \bar{\gamma} f_{i,j-1} + f_{i-1,j-1} && \text{for } i = s - q + 2, \dots, s, \quad j = 2, \dots, t, \end{aligned}$$

and  $f_{i,1}$  for  $i = s - q + 2, \dots, s$  is defined as the solution to

$$\langle F_{s \times t}[q]G_{t-1}^T - \lambda F_{s \times t}[q]\hat{G}_{t-1}^T, F_{s \times t}[s - i + 1]G_{t-1}^T - \lambda F_{s \times t}[s - i + 1]\hat{G}_{t-1}^T \rangle \equiv 0.$$

Notice that  $f_{i,1}$  is used as an unknown in the generation of elements in (5.5). In the definition of  $F_{s \times t}[q]$ , the solutions for  $f_{i,1}$  for  $i = s - q + 2, \dots, s$  ensure that  $F_{s \times t}[q]G_{t-1}^T - \lambda F_{s \times t}[q]\hat{G}_{t-1}^T$  is orthogonal to  $F_{s \times t}[\hat{q}]G_{t-1}^T - \lambda F_{s \times t}[\hat{q}]\hat{G}_{t-1}^T$  for  $\hat{q} = 1, \dots, q - 1$ .

Also here we show a small example to facilitate the interpretation of the definition:

$$F_{3 \times 2}^D = \begin{bmatrix} p_1 & p_1 \bar{\gamma} \\ p_2 - p_1 \frac{(|\gamma|^2 + 1)\gamma}{|\gamma|^4 + 2|\gamma|^2 + 2} & p_2 \bar{\gamma} + p_1 \frac{|\gamma|^2 + 2}{|\gamma|^4 + 2|\gamma|^2 + 2} \\ p_3 - p_2 \frac{\bar{\gamma}}{|\gamma|^2 + 1} + p_1 \frac{\bar{\gamma}^2}{|\gamma|^4 + 2|\gamma|^2 + 2} & p_3 \bar{\gamma} + p_2 \frac{1}{|\gamma|^2 + 1} - p_1 \frac{\bar{\gamma}}{|\gamma|^4 + 2|\gamma|^2 + 2} \end{bmatrix}.$$

**5.3. Versal deformations—the general case.** Without loss of generality assume that  $A - \lambda B$  is already in KCF,  $M = \text{diag}(M_1, M_2, \dots, M_b)$ , where each  $M_k$  is either a Jordan block associated with a finite or infinite eigenvalue or a singular block corresponding to a left or right minimal index. A pencil  $T_A - \lambda T_B = XM - MY$  in the tangent space can be partitioned conformally with the pencil  $M$  so that

$T_{ij}^A - \lambda T_{ij}^B = X_{ij}M_j - M_iY_{ij}$ , where  $M_k$  is  $m_k \times n_k$ ,  $X_{ij}$  is  $m_i \times m_j$ , and  $Y_{ij}$  is  $n_i \times n_j$ :

$$\begin{bmatrix} X_{11} & \cdots & X_{1b} \\ \vdots & \ddots & \vdots \\ X_{b1} & \cdots & X_{bb} \end{bmatrix} \begin{bmatrix} M_1 & & \\ & \ddots & \\ & & M_b \end{bmatrix} - \begin{bmatrix} M_1 & & \\ & \ddots & \\ & & M_b \end{bmatrix} \begin{bmatrix} Y_{11} & \cdots & Y_{1b} \\ \vdots & \ddots & \vdots \\ Y_{b1} & \cdots & Y_{bb} \end{bmatrix}.$$

Since the blocks  $T_{ij}^A - \lambda T_{ij}^B, i, j = 1, \dots, b$  are mutually independent, we can study the different blocks of  $T_A - \lambda T_B$  separately. Let  $Z_{ij}^A - \lambda Z_{ij}^B$  be conformally sized blocks of  $Z_A - \lambda Z_B$ . From (4.4) we know that  $Z_A - \lambda Z_B$  is in the normal space if and only if  $A^H Z_A + B^H Z_B = 0$  and  $Z_A A^H + Z_B B^H = 0$ . We obtain a simple result since  $A$  and  $B$  are block diagonal.

PROPOSITION 5.2. *Assume that*

$$M = A - \lambda B = \text{diag}(A_1, A_2, \dots, A_b) - \lambda \text{diag}(B_1, B_2, \dots, B_b)$$

*is in KCF, where each block  $A_i - \lambda B_i \equiv M_i$  represents one block in the Kronecker structure. Then  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$  if and only if*

$$A_j^H Z_{ji}^A = -B_j^H Z_{ji}^B \quad \text{and} \quad Z_{ji}^A A_i^H = -Z_{ji}^B B_i^H \quad \text{for } i = 1, \dots, b \text{ and } j = 1, \dots, b.$$

The mutual independency of the  $(i, j)$  blocks of  $Z_A$  and  $Z_B$  implies that we only have to consider two  $M_k$  blocks at a time:

$$\begin{aligned} T_A[i, j] - \lambda T_B[i, j] &= \begin{bmatrix} X_{ii} & X_{ij} \\ X_{ji} & X_{jj} \end{bmatrix} \begin{bmatrix} M_i & 0 \\ 0 & M_j \end{bmatrix} - \begin{bmatrix} M_i & 0 \\ 0 & M_j \end{bmatrix} \begin{bmatrix} Y_{ii} & Y_{ij} \\ Y_{ji} & Y_{jj} \end{bmatrix} \\ &= \begin{bmatrix} T_{ii}^A & T_{ij}^A \\ T_{ji}^A & T_{jj}^A \end{bmatrix} - \lambda \begin{bmatrix} T_{ii}^B & T_{ij}^B \\ T_{ji}^B & T_{jj}^B \end{bmatrix} \end{aligned}$$

and

$$(5.6) \quad Z_A[i, j] - \lambda Z_B[i, j] = \begin{bmatrix} Z_{ii}^A & Z_{ij}^A \\ Z_{ji}^A & Z_{jj}^A \end{bmatrix} - \lambda \begin{bmatrix} Z_{ii}^B & Z_{ij}^B \\ Z_{ji}^B & Z_{jj}^B \end{bmatrix}.$$

Notably, by interchanging the blocks  $M_i = A_i - \lambda B_i$  and  $M_j = A_j - \lambda B_j$  in the KCF, we only have to interchange the corresponding blocks in  $Z_A - \lambda Z_B$  accordingly. For example, if  $Z_A[i, j] - \lambda Z_B[i, j]$  in (5.6) belongs to  $\text{nor}(\text{diag}(M_i, M_j))$ , then

$$\begin{bmatrix} Z_{jj}^A & Z_{ji}^A \\ Z_{ij}^A & Z_{ii}^A \end{bmatrix} - \lambda \begin{bmatrix} Z_{jj}^B & Z_{ji}^B \\ Z_{ij}^B & Z_{ii}^B \end{bmatrix} \in \text{nor}(\text{diag}(M_j, M_i)).$$

This implies that given two blocks  $M_i$  and  $M_j$ , it is sufficient to consider the case  $\text{diag}(M_i, M_j)$ . In the following we will order the blocks in the KCF so that  $Z_A - \lambda Z_B$  is block lower triangular.

THEOREM 5.3. *Let  $A - \lambda B = \text{diag}(A_1, A_2, \dots, A_b) - \lambda \text{diag}(B_1, B_2, \dots, B_b)$  be in KCF with the structure blocks  $M_i = A_i - \lambda B_i$  ordered as follows:  $L_k, J_k(0), J_k(\gamma)$  (for  $\gamma \neq \{0, \infty\}$ ),  $N_k$ , and  $L_k^T$ , where the ordering within each block type is in increasing order of size, except for the  $L_k^T$  blocks, which are ordered by decreasing order of size.*

*For all  $i$  and  $j$ , let the  $(i, j), (j, i)$  and  $(i, i), (j, j)$  blocks of  $Z_A(p) - \lambda Z_B(p)$  corresponding to  $\text{diag}(M_i, M_j)$  be built from Table 5.1 and Table 5.2, respectively.*

TABLE 5.1

Blocks in  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$ , where for  $L_\alpha \oplus L_\beta$ ,  $J_\alpha(0) \oplus J_\beta(0)$ ,  $J_\alpha(\gamma) \oplus J_\beta(\gamma)$ , and  $N_\alpha \oplus N_\beta$  it is assumed that  $\alpha \leq \beta$ . For  $L_\alpha^T \oplus L_\beta^T$ ,  $\alpha \geq \beta$  is assumed. Also  $\gamma_1 \neq \gamma_2$  is assumed.

KCF: $M_i \oplus M_j$	$Z_{ij}^A$	$Z_{ij}^B$	$Z_{ji}^A$	$Z_{ji}^B$
$L_\alpha \oplus L_\beta$	0	0	$S_{\beta \times (\alpha+1)}^B$	$-C_\beta^T S_{\beta \times (\alpha+1)}^B$
$L_\alpha \oplus J_\beta(0)$	0	0	$S_{\beta \times (\alpha+1)}^L$	$-C_\beta^T S_{\beta \times (\alpha+1)}^L$
$L_\alpha \oplus J_\beta(\gamma)$	0	0	$F_{\beta \times (\alpha+2)}^D G_{\alpha+1}^T$	$-F_{\beta \times (\alpha+2)}^D \hat{G}_{\alpha+1}^T$
$L_\alpha \oplus N_\beta$	0	0	$C_\beta^T H_{\beta \times (\alpha+1)}^L$	$-H_{\beta \times (\alpha+1)}^L$
$L_\alpha \oplus L_\beta^T$	0	0	$G_{\beta+1} H_{(\beta+2) \times (\alpha+1)}$	$-\hat{G}_{\beta+1} H_{(\beta+2) \times (\alpha+1)}$
$J_\alpha(0) \oplus J_\beta(0)$	$S_{\alpha \times \beta}^L$	$-C_\alpha^T S_{\alpha \times \beta}^L$	$T_{\beta \times \alpha}^L$	$-C_\beta^T T_{\beta \times \alpha}^L$
$J_\alpha(0) \oplus L_\beta^T$	0	0	$H_{(\beta+1) \times \alpha}^U$	$-H_{(\beta+1) \times \alpha}^U C_\alpha^T$
$J_\alpha(\gamma) \oplus J_\beta(\gamma)$	$D_{\alpha \times \beta}^L$	$E_{\alpha \times \beta}^L$	$D_{\beta \times \alpha}^L$	$E_{\beta \times \alpha}^L$
$J_\alpha(\gamma) \oplus L_\beta^T$	0	0	$G_{\beta+1} (\hat{I}_\alpha F_{\alpha \times (\beta+2)}^D)^T$	$-\hat{G}_{\beta+1} (\hat{I}_\alpha F_{\alpha \times (\beta+2)}^D)^T$
$N_\alpha \oplus N_\beta$	$C_\alpha^T S_{\alpha \times \beta}^L$	$-S_{\alpha \times \beta}^L$	$C_\beta^T T_{\beta \times \alpha}^L$	$-T_{\beta \times \alpha}^L$
$N_\alpha \oplus L_\beta^T$	0	0	$T_{(\beta+1) \times \alpha}^L C_\alpha^T$	$-T_{(\beta+1) \times \alpha}^L$
$L_\alpha^T \oplus L_\beta^T$	0	0	$T_{(\beta+1) \times \alpha}^B$	$-T_{(\beta+1) \times \alpha}^B C_\alpha$
$J_\alpha(0) \oplus J_\beta(\gamma)$	0	0	0	0
$J_\alpha(0) \oplus N_\beta$	0	0	0	0
$J_\alpha(\gamma_1) \oplus J_\beta(\gamma_2)$	0	0	0	0
$J_\alpha(\gamma) \oplus N_\beta$	0	0	0	0

TABLE 5.2

The diagonal blocks in  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$ .

KCF: $M_i$	$Z_{ii}^A$	$Z_{ii}^B$
$L_\alpha$	0	0
$J_\alpha(0)$	$S_{\alpha \times \alpha}^L$	$-C_\alpha^T S_{\alpha \times \alpha}^L$
$J_\alpha(\gamma)$	$D_{\alpha \times \alpha}^L$	$E_{\alpha \times \alpha}^L$
$N_\alpha$	$C_\alpha^T S_{\alpha \times \alpha}^L$	$-S_{\alpha \times \alpha}^L$
$L_\alpha^T$	0	0

Then  $Z_A(p) - \lambda Z_B(p)$  gives an orthogonal basis for  $\text{nor}(A - \lambda B)$  with minimum number of parameters; i.e.,  $\mathcal{V}(p) = A - \lambda B + Z_A(p) - \lambda Z_B(p)$  is a miniversal deformation of  $A - \lambda B$ .

The superscripts  $B, L, U$ , and  $D$  of the matrices in Tables 5.1 and 5.2 are parts of the matrix definitions in section 5.2. The superscript  $T$  is the matrix transpose. All subscripts, e.g.,  $\alpha \times \beta$ , refer to the sizes of the matrices.

Notice that the diagonal blocks  $(i, i)$  and  $(j, j)$  of  $Z_A - \lambda Z_B$  can also be obtained from Table 5.1 by setting  $i = j$ . For clarity we also display the expressions for the  $(i, i)$  and  $(j, j)$  blocks of  $Z_A - \lambda Z_B$  corresponding to all kinds of structure blocks  $M_i$  in Table 5.2. Of course, the  $(j, j)$  blocks corresponding to  $M_j$  are read from Table 5.2 by substituting  $\alpha$  with  $\beta$ .

The proof of Theorem 5.3 consists of three parts.

1. The blocks of  $Z_A - \lambda Z_B$  displayed in Table 5.1 fulfill the conditions in Proposition 5.2, which imply that  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$  is orthogonal to an arbitrary  $T_A - \lambda T_B \in \text{tan}(A - \lambda B)$ .
2. The number of independent parameters in  $Z_A - \lambda Z_B$  is equal to the codimension of  $\text{orbit}(A - \lambda B)$ , which implies that the parameterized normal form has minimum number of parameters.
3. Each block in Table 5.1 defines an orthogonal basis; i.e., the basis for each parameter  $p_i$  is orthogonal to the basis for each other parameter  $p_j$ ,  $i \neq j$ .

We start by proving part 3 and then prove parts 1 and 2 for the 16 different cases  $\text{diag}(M_i, M_j)$  corresponding to different combinations of structure blocks in the KCF. In Table 5.3 we display the codimension for these 16 cases and the number of parameters in the  $(i, i)$ ,  $(i, j)$ ,  $(j, i)$ , and  $(j, j)$  blocks of  $Z_A - \lambda Z_B$ . The codimensions are computed from (4.8), which is the minimum number of parameters required to span the corresponding normal space. For the ordering and the sizes of the blocks in  $A - \lambda B$  we have made the same assumptions in Table 5.3 as in Table 5.1. Notice that the codimension counts for  $L_\alpha \oplus L_\beta$  and  $L_\alpha^T \oplus L_\beta^T$  are 0 if  $\alpha = \beta$ . The number of parameters required in each of the  $(i, i)$ ,  $(i, j)$ ,  $(j, i)$ , and  $(j, j)$  blocks of  $Z_A - \lambda Z_B$  follows from the proof given below.

TABLE 5.3

The number of parameters in the  $(i, i)$ ,  $(i, j)$ ,  $(j, i)$ , and  $(j, j)$  blocks of  $Z_A - \lambda Z_B \in \text{nor}(M_i \oplus M_j)$ .

KCF: $M_i \oplus M_j$	$\text{cod}(M_i \oplus M_j)$	$(i, i)$	$(i, j)$	$(j, i)$	$(j, j)$
$L_\alpha \oplus L_\beta$	$\beta - \alpha - 1$	0	0	$\beta - \alpha - 1$	0
$L_\alpha \oplus J_\beta(0)$	$2\beta$	0	0	$\beta$	$\beta$
$L_\alpha \oplus J_\beta(\gamma)$	$2\beta$	0	0	$\beta$	$\beta$
$L_\alpha \oplus N_\beta$	$2\beta$	0	0	$\beta$	$\beta$
$L_\alpha \oplus L_\beta^T$	$\alpha + \beta + 2$	0	0	$\alpha + \beta + 2$	0
$J_\alpha(0) \oplus J_\beta(0)$	$\beta + 3\alpha$	$\alpha$	$\alpha$	$\alpha$	$\beta$
$J_\alpha(0) \oplus L_\beta^T$	$2\alpha$	$\alpha$	0	$\alpha$	0
$J_\alpha(\gamma) \oplus J_\beta(\gamma)$	$\beta + 3\alpha$	$\alpha$	$\alpha$	$\alpha$	$\beta$
$J_\alpha(\gamma) \oplus L_\beta^T$	$2\alpha$	$\alpha$	0	$\alpha$	0
$N_\alpha \oplus N_\beta$	$\beta + 3\alpha$	$\alpha$	$\alpha$	$\alpha$	$\beta$
$N_\alpha \oplus L_\beta^T$	$2\alpha$	$\alpha$	0	$\alpha$	0
$L_\alpha^T \oplus L_\beta^T$	$\alpha - \beta - 1$	0	0	$\alpha - \beta - 1$	0
$J_\alpha(0) \oplus J_\beta(\gamma)$	$\alpha + \beta$	$\alpha$	0	0	$\beta$
$J_\alpha(0) \oplus N_\beta$	$\alpha + \beta$	$\alpha$	0	0	$\beta$
$J_\alpha(\gamma_1) \oplus J_\beta(\gamma_2)$	$\alpha + \beta$	$\alpha$	0	0	$\beta$
$J_\alpha(\gamma) \oplus N_\beta$	$\alpha + \beta$	$\alpha$	0	0	$\beta$

To fully appreciate this rather technical proof it could be more fruitful to look first at some examples of versal deformations in section 6.1.

*Proof of part 3.* We show that each matrix pencil block in Table 5.1 has all its parameters in orthogonal directions. This is trivial for blocks built from the structured Toeplitz and Hankel matrices  $S^L$ ,  $S^B$ ,  $H$ ,  $H^L$ ,  $H^U$ ,  $T^L$ , or  $T^B$  (possibly involving some kind of shift). Remember that the Frobenius inner product can be expressed in terms of the sum of all results from elementwise multiplications as shown in (5.1). For each of these matrices, the elementwise multiplication of the basis for one parameter  $p_i$  and the basis for another parameter  $p_j$ ,  $j \neq i$  only results in multiplications where at least one of the two elements is zero. Obviously, these bases are orthogonal. For the matrix pencil blocks built from the  $F^D$  matrix, the orthogonality follows from

construction since some of the elements are explicitly chosen so that the Frobenius inner product is zero.

For the proof for the blocks of type  $D^L - \lambda E^L$  we define  $s_q$  in terms of the  $d_i$  and  $e_i$  in (5.3) to be

$$s_q = \sum_{i=1}^q i|d_i|^2 + \sum_{i=1}^{q-1} i|e_i|^2 - q\bar{\gamma}d_q\bar{e}_q.$$

Independent of  $s$  and  $t$ , the number  $s_q$  is the inner product of the  $q$ th basis vector with the  $r$ th, where  $q < r$ .

We show by induction that  $s_q = 0$  for  $q = 1, 2, \dots$ . Clearly  $s_1 = |\gamma|^2 - \gamma\bar{\gamma} = 0$ .

We now show that  $s_{q+1} - s_q = 0$ , from which the result follows:

$$\begin{aligned} & q\bar{\gamma}d_q\bar{e}_q + (q+1)|d_{q+1}|^2 + q|e_q|^2 - (q+1)\bar{\gamma}d_{q+1}\bar{e}_{q+1} \\ &= q\bar{e}_q(\bar{\gamma}d_q + e_q) + (q+1)d_{q+1}(\bar{d}_{q+1} - \bar{\gamma}\bar{e}_{q+1}) \\ &= d_{q+1}((q+1)(\bar{d}_{q+1} - \bar{\gamma}\bar{e}_{q+1}) - q\bar{e}_q) \\ &= d_{q+1}\left((q+1)\left(-\gamma\bar{d}_q - \bar{e}_q + \gamma\bar{d}_q + 2\bar{e}_q - \frac{\bar{e}_q}{q+1}\right) - q\bar{e}_q\right) \\ &= d_{q+1}((q+1)\bar{e}_q - \bar{e}_q - q\bar{e}_q) = 0. \end{aligned}$$

Since  $Z_A - \lambda Z_B$  is built from  $b^2$  mutually independent blocks in Table 5.1, each associated with  $c_i$  parameters, it follows that  $Z_A - \lambda Z_B$  is an orthogonal basis for a  $(c_1 + c_2 + \dots + c_{b^2})$ -dimensional space, with one parameter for each dimension.  $\square$

*Proof of parts 1 and 2.* Now, it remains to show that  $Z_A - \lambda Z_B$  is orthogonal to  $\tan(A - \lambda B)$  and that the number of parameters in  $Z_A - \lambda Z_B$  is equal to  $\text{cod}(A - \lambda B)$ . Since the number of parameters in orthogonal directions cannot exceed the codimension, it is sufficient to show that we have found them all. The orthogonality between  $Z_A - \lambda Z_B$  and  $\tan(A - \lambda B)$  is shown by proving that each pair of blocks fulfills the conditions  $A_j^H Z_{ji}^A = -B_j^H Z_{ji}^B$  and  $Z_{ji}^A A_i^H = -Z_{ji}^B B_i^H$  in Proposition 5.2. In the following we refer to these as the *first* and *second* conditions, respectively.

We carry out the proofs for all 16 cases  $M_i \oplus M_j$  in Table 5.1, starting with blocks where  $M_i$  and  $M_j$  are of the same kind.

$\mathbf{J}_\alpha(\mathbf{0}) \oplus \mathbf{J}_\beta(\mathbf{0})$ : We note that  $J_k(0) = C_k - \lambda I_k$ . *First* condition for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = C_\beta^T T_{\beta \times \alpha}^L = I_\beta C_\beta^T T_{\beta \times \alpha}^L = -B_j^H Z_{ji}^B.$$

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H = T_{\beta \times \alpha}^L C_\alpha^T = T_{\beta \times \alpha}^L C_\alpha^T I_\alpha = C_\beta^T T_{\beta \times \alpha}^L I_\alpha = -Z_{ji}^B B_i^H,$$

where we used that  $T_{\beta \times \alpha}^L C_\alpha^T = C_\beta^T T_{\beta \times \alpha}^L$  for  $\beta \geq \alpha$ . Similarly for the  $(i, j)$  block,

$$A_i^H Z_{ij}^A = C_\alpha^T S_{\alpha \times \beta}^L = I_\alpha C_\alpha^T S_{\alpha \times \beta}^L = -B_i^H Z_{ij}^B$$

and

$$Z_{ij}^A A_j^H = S_{\alpha \times \beta}^L C_\beta^T = S_{\alpha \times \beta}^L C_\beta^T I_\beta = C_\alpha^T S_{\alpha \times \beta}^L I_\beta = -Z_{ij}^B B_j^H.$$

Here we used that  $S_{\alpha \times \beta}^L C_\beta^T = C_\alpha^T S_{\alpha \times \beta}^L$  for  $\beta \geq \alpha$ .

Since the  $(i, i)$ ,  $(i, j)$ , and  $(j, i)$  blocks of  $Z_A - \lambda Z_B$  have  $\alpha$  parameters each and the  $(j, j)$  block has  $\beta$  parameters, the total number of parameters in  $Z_A - \lambda Z_B$  is equal to  $\text{cod}(J_\alpha(0) \oplus J_\beta(0)) = \beta + 3\alpha$ .

$\mathbf{N}_\alpha \oplus \mathbf{N}_\beta$ : Since there is a symmetry between  $J_k(0) = C_k - \lambda I_k$  and  $N_k = I_k - \lambda C_k$  and there is a corresponding symmetry between blocks in  $Z_A - \lambda Z_B$  for  $J_k(0)$  and  $N_k$  blocks, the proof for  $N_\alpha \oplus N_\beta$  is similar to the case  $J_\alpha(0) \oplus J_\beta(0)$ .

$\mathbf{J}_\alpha(\gamma) \oplus \mathbf{J}_\beta(\gamma)$ : Here the  $(j, i)$  block and the  $(i, j)$  block are defined similarly (see Table 5.1), and therefore it is sufficient to prove one of them with no constraints on  $\alpha$  and  $\beta$ . We note that  $J_k(\gamma) = \gamma I_k + C_k - \lambda I_k$ . We show that the first and second conditions hold for  $Z_{ji}^A = D_{\beta \times \alpha}[q]$  and  $Z_{ji}^B = E_{\beta \times \alpha}[q]$  for  $q = 1, \dots, \min\{\alpha, \beta\}$ . *First condition*:

$$A_j^H Z_{ji}^A = (\gamma I_\beta + C_\beta)^H D_{\beta \times \alpha}[q] = \bar{\gamma} D_{\beta \times \alpha}[q] + C_\beta^T D_{\beta \times \alpha}[q].$$

Remember that  $D_{\beta \times \alpha}[q]$  has all elements zero, except for the  $q$  lower left diagonals, where all elements in each diagonal are identical and defined by the element in the first column. For  $q = 1$  the proof is trivial. For  $q > 1$ ,  $A_j^H Z_{ji}^A$  gives the following matrix. All diagonals starting at position  $(u, 1)$  for  $1 \leq u \leq \beta - q$  are zero. The elements in the diagonal starting at position  $(\beta - q + 1, 1)$  are  $\bar{\gamma} d_q$ , which by definition is equal to  $-e_q^*$ , which in turn defines the corresponding diagonal in  $-E_{\beta \times \alpha}[q]$ . The elements in the diagonals starting at positions  $(\beta - u + 1, 1)$ , where  $1 \leq u < q$ , are equal to  $\bar{\gamma} d_u + d_{u+1}$ . Since  $d_{u+1}$  is defined as  $-\bar{\gamma} d_u - e_u$ , the elements in these diagonals are equal to  $-e_u$ , which defines the elements in the corresponding diagonals in  $-E_{\beta \times \alpha}[q]$ . Since  $-E_{\beta \times \alpha}[q] = -B_j^H Z_{ji}^B$ , we have proved the first condition.

*Second condition*: Since  $D_{\beta \times \alpha}[q]$  only has  $q \leq \min\{s, t\}$  nonzero diagonals in the lower left corner of the matrix, a shift of rows downward gives the same result as a shift of columns leftward, i.e.,  $C_\beta^T D_{\beta \times \alpha}[q] = D_{\beta \times \alpha}[q] C_\alpha^T$ . Using information from the first part, we obtain

$$\begin{aligned} Z_{ji}^A A_i^H &= D_{\beta \times \alpha}[q] (\gamma I_\alpha + C_\alpha)^H = \bar{\gamma} D_{\beta \times \alpha}[q] + D_{\beta \times \alpha}[q] C_\alpha^T = \bar{\gamma} D_{\beta \times \alpha}[q] + C_\beta^T D_{\beta \times \alpha}[q] \\ &= A_j^H Z_{ji}^A = -E_{\beta \times \alpha}[q] = -Z_{ji}^B B_i^H \end{aligned}$$

since  $B_i$  is the identity matrix.

Also here, the number of parameters in  $Z_{jj}^A - \lambda Z_{jj}^B$  is  $\beta$ , and there are  $\alpha$  parameters in each of the other three blocks, giving  $\beta + 3\alpha$  in total.

Even though the  $(i, i)$ ,  $(j, i)$ ,  $(i, j)$ , and  $(j, j)$  blocks look rather complicated, they reduce for  $\gamma = 0$  to the corresponding blocks for  $J_\alpha(0) \oplus J_\beta(0)$  in Table 5.1.

$\mathbf{L}_\alpha \oplus \mathbf{L}_\beta$ : Here we use  $L_k = \hat{G}_k - \lambda G_k$ . *First condition* for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = \hat{G}_\beta^T S_{\beta \times (\alpha+1)}^B = \begin{bmatrix} 0 \\ S_{\beta \times (\alpha+1)}^B \end{bmatrix} = \begin{bmatrix} C_\beta^T S_{\beta \times (\alpha+1)}^B \\ 0 \end{bmatrix} = G_\beta^T C_\beta^T S_{\beta \times (\alpha+1)}^B = -B_j^H Z_{ji}^B.$$

*Second condition* for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H = S_{\beta \times (\alpha+1)}^B \hat{G}_\beta^T = \begin{bmatrix} 0 \\ S_{\beta \times (\alpha+1)}^B \end{bmatrix} = \begin{bmatrix} C_\beta^T S_{\beta \times (\alpha+1)}^B \\ 0 \end{bmatrix} = C_\beta^T S_{\beta \times (\alpha+1)}^B G_\beta^T = -Z_{ji}^B B_i^H.$$

Since the contribution from  $L_\alpha \oplus L_\beta$  to the codimension is  $\beta - \alpha - 1$  and the  $(j, i)$  block has  $\beta - \alpha - 1$  independent parameters, we deduce that all other blocks in  $Z_A - \lambda Z_B$  are zero.

$\mathbf{L}_\alpha^T \oplus \mathbf{L}_\beta^T$ : Since this case is just the transpose of  $L_\alpha \oplus L_\beta$ , the proof is almost the same, and therefore we omit the technical details here.

So far we have proved all cases where both blocks are of the same type. Since the diagonal blocks in  $Z_A - \lambda Z_B$  always correspond to such cases (see Table 5.3 for the number of parameters in these blocks), from now on we only have to consider the  $(i, j)$  and  $(j, i)$  blocks, where  $i \neq j$  for the remaining cases.

$\mathbf{L}_\alpha \oplus \mathbf{J}_\beta(\mathbf{0})$ : *First* condition for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = C_\beta^T S_{\beta \times (\alpha+1)}^L = I_\beta C_\beta^T S_{\beta \times (\alpha+1)}^L = -B_j^H Z_{ji}^B.$$

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H = S_{\beta \times (\alpha+1)}^L \hat{G}_\alpha^T = C_\beta^T S_{\beta \times (\alpha+1)}^L G_\alpha^T = -Z_{ji}^B B_i^H.$$

The  $(i, i)$  and  $(j, j)$  blocks contribute with zero and  $\beta$  parameters, respectively. Since the  $(j, i)$  block gives another  $\beta$  parameters, we have found all  $2\beta$  parameters, and therefore it follows that  $Z_{ij}^A = \lambda Z_{ij}^B = 0$ .

$\mathbf{L}_\alpha \oplus \mathbf{J}_\beta(\gamma)$ : *First* condition for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = (\gamma I_\beta + C_\beta)^H F_{\beta \times (\beta+2)}^D G_{\alpha+1}^T = \bar{\gamma} F_{\beta \times (\beta+2)}^D G_{\alpha+1}^T + C_\beta^T F_{\beta \times (\beta+2)}^D G_{\alpha+1}^T.$$

By inspection we see that the  $(u, v)$  element of this matrix is  $\bar{\gamma} f_{u,v}^d + f_{u-1,v}^d$  if  $u > 1$  and  $\bar{\gamma} f_{u,v}^d$  if  $u = 1$  (where  $f_{u,v}^d$  denotes the  $(u, v)$  element of  $F^D$ ). The right-hand side of the same condition is

$$-B_j^H Z_{ji}^B = I_\beta F_{\beta \times (\beta+2)}^D \hat{G}_{\alpha+1}^T,$$

which simply is the  $\beta$  leftmost columns of  $F_{\beta \times (\beta+2)}^D$ . The  $(u, v)$  element of this matrix is then  $f_{u,v+1}^d$ , which is defined as  $\bar{\gamma} f_{u,v}^d + f_{u-1,v}^d$  if  $u > 1$  and  $\bar{\gamma} f_{u,v}^d$  if  $u = 1$ .

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H \hat{G}_\alpha^T = F_{\beta \times (\alpha+2)}^D G_{\alpha+1}^T \hat{G}_\alpha^T = F_{\beta \times (\alpha+2)}^D \begin{bmatrix} 0 \\ I_\alpha \\ 0 \end{bmatrix} = F_{\beta \times (\alpha+2)}^D \hat{G}_{\alpha+1} G_\alpha^T = -Z_{ji}^B B_i^H.$$

As in the previous case, the  $(i, i)$  and  $(j, j)$  blocks contribute with zero and  $\beta$  parameters, respectively. Since the  $(j, i)$  block gives the remaining  $\beta$  parameters, the  $(i, j)$  block is the zero pencil.

Notably, for  $\gamma = 0$ , the “monstrous”  $(j, i)$  block reduces to the  $(j, i)$  block for  $L_\alpha \oplus J_\beta(0)$  in Table 5.1.

$\mathbf{L}_\alpha \oplus \mathbf{N}_\beta$ : *First* condition for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = I_\beta C_\beta^T H_{\beta \times (\alpha+1)}^L = C_\beta^T H_{\beta \times (\alpha+1)}^L = -B_j^H Z_{ji}^B.$$

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H \hat{G}_\alpha^T = C_\beta^T H_{\beta \times (\alpha+1)}^L = \begin{bmatrix} 0 \\ H_{(\beta-1) \times \alpha}^L \end{bmatrix} = H_{\beta \times (\alpha+1)}^L G_\alpha^T = -Z_{ji}^B B_i^H.$$

Also here, the  $(i, i)$  and  $(j, j)$  blocks contribute with zero and  $\beta$  parameters, respectively. Since the  $(j, i)$  block gives the remaining  $\beta$  parameters, the  $(i, j)$  block is the zero pencil.

$\mathbf{L}_\alpha \oplus \mathbf{L}_\beta^{\mathbf{T}}$ : For this case the  $(i, i)$  and  $(j, j)$  blocks are zero pencils. *First* condition for the  $(j, i)$  block:

$$\begin{aligned} A_j^H Z_{ji}^A &= \hat{G}_\beta G_{\beta+1} H_{(\beta+2) \times (\alpha+1)} = [0 \ I_\beta \ 0] H_{(\beta+2) \times (\alpha+1)} \\ &= G_\beta \hat{G}_{\beta+1} H_{(\beta+2) \times (\alpha+1)} = -B_j^H Z_{ji}^B. \end{aligned}$$

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H = G_{\beta+1} H_{(\beta+2) \times (\alpha+1)} \hat{G}_\alpha^T,$$

which is a matrix consisting of the  $\beta+1$  first rows and  $\alpha$  last columns of  $H_{(\beta+2) \times (\alpha+1)}$ . This matrix is identical to the one given by the  $\beta+1$  last rows and  $\alpha$  first columns of  $H_{(\beta+2) \times (\alpha+1)}$ , i.e.,

$$\hat{G}_{\beta+1} H_{(\beta+2) \times (\alpha+1)} G_\alpha^T = -Z_{ji}^B B_i^H.$$

Since this block has all  $\alpha + \beta + 2$  parameters, it follows that the  $(i, j)$  block is the zero pencil.

$\mathbf{J}_\alpha(\mathbf{0}) \oplus \mathbf{L}_\beta^{\mathbf{T}}$ : *First* condition for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = \hat{G}_\beta H_{(\beta+1) \times \alpha}^U,$$

which simply is the last  $\beta$  rows in  $H_{(\beta+1) \times \alpha}^U$ . Another way to construct this matrix is to shift the columns in  $H_{(\beta+1) \times \alpha}^U$  one column leftward and pick the  $\beta$  first columns of the matrix, which can be written as

$$G_\beta H_{(\beta+1) \times \alpha}^U C_\alpha^T = -B_j^H Z_{ji}^B.$$

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H = H_{(\beta+1) \times \alpha}^U C_\alpha^T = H_{(\beta+1) \times \alpha}^U C_\alpha^T I_\alpha = -Z_{ji}^B B_i^H.$$

The  $(i, i)$  and  $(j, j)$  blocks contribute with  $\alpha$  and zero parameters, respectively. Since the  $(j, i)$  block gives another  $\alpha$  parameters, we conclude that the  $(i, j)$  block is the zero pencil.

$\mathbf{J}_\alpha(\gamma) \oplus \mathbf{L}_\beta^{\mathbf{T}}$ : Since the proof for this case is similar to the one for the case  $L_\alpha \oplus J_\beta(\gamma)$ , we omit the technical details here. It follows that for  $\gamma = 0$ , the  $(j, i)$  block reduces to the  $(j, i)$  block for  $J_\alpha(0) \oplus L_\beta^{\mathbf{T}}$  in Table 5.1.

$\mathbf{N}_\alpha \oplus \mathbf{L}_\beta^{\mathbf{T}}$ : *First* condition for the  $(j, i)$  block:

$$A_j^H Z_{ji}^A = \hat{G}_\beta T_{(\beta+1) \times \alpha}^L C_\alpha^T,$$

which is the last  $\beta$  rows in  $T_{(\beta+1) \times \alpha}^L$  shifted one column leftward. This matrix is identical to the one given by the  $\beta$  first rows in  $T_{(\beta+1) \times \alpha}^L$ , which is

$$G_\beta T_{(\beta+1) \times \alpha}^L = -B_j^H Z_{ji}^B.$$

*Second* condition for the  $(j, i)$  block:

$$Z_{ji}^A A_i^H = T_{(\beta+1) \times \alpha}^L C_\alpha^T I_\alpha = T_{(\beta+1) \times \alpha}^L C_\alpha^T = -Z_{ji}^B B_i^H.$$



The  $(i, i)$  and  $(j, j)$  blocks in  $Z_A - \lambda Z_B$  contribute with  $\alpha$  and zero parameters, respectively. Since the  $(j, i)$  block gives another  $\alpha$  parameters, we conclude that the  $(i, j)$  block is the zero pencil.

$\mathbf{J}_\alpha(\mathbf{0}) \oplus \mathbf{J}_\beta(\gamma)$ ,  $\mathbf{J}_\alpha(\mathbf{0}) \oplus \mathbf{N}_\beta$ ,  $\mathbf{J}_\alpha(\gamma_1) \oplus \mathbf{J}_\beta(\gamma_2)$ , and  $\mathbf{J}_\alpha(\gamma) \oplus \mathbf{N}_\beta$ : In these four cases the  $(i, i)$  and  $(j, j)$  blocks contribute with  $\alpha$  and  $\beta$  parameters, respectively, and therefore the  $(j, i)$  and  $(i, j)$  blocks are zero pencils.

Since we have considered all possible cases of  $M_i$  and  $M_j$  blocks, the proof is complete.  $\square$

**6. Applications and examples.**

**6.1. Some examples of versal deformations of matrix pencils in KCF.**

In the following we show three examples of versal deformations of matrix pencils. For the  $7 \times 8$  pencil  $A - \lambda B = L_2 \oplus J_2(0) \oplus J_3(0)$  with codimension 14, the 14-parameter versal deformation  $\mathcal{V}(p) = A - \lambda B + Z_A - \lambda Z_B$ , where  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$ , is given by

$$Z_A = \left[ \begin{array}{ccc|ccc|ccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline p_1 & 0 & 0 & p_6 & 0 & p_{10} & 0 & 0 & 0 \\ p_2 & p_1 & 0 & p_7 & p_6 & p_{11} & p_{10} & 0 & 0 \\ \hline p_3 & 0 & 0 & 0 & 0 & p_{12} & 0 & 0 & 0 \\ p_4 & p_3 & 0 & p_8 & 0 & p_{13} & p_{12} & 0 & 0 \\ p_5 & p_4 & p_3 & p_9 & p_8 & p_{14} & p_{13} & p_{12} & 0 \end{array} \right]$$

and

$$Z_B = \left[ \begin{array}{ccc|ccc|ccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -p_1 & 0 & 0 & -p_6 & 0 & -p_{10} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -p_3 & 0 & 0 & 0 & 0 & -p_{12} & 0 & 0 & 0 \\ -p_4 & -p_3 & 0 & -p_8 & 0 & -p_{13} & -p_{12} & 0 & 0 \end{array} \right].$$

For the  $3 \times 4$  pencil  $A - \lambda B = L_1 \oplus J_2(\gamma)$  with codimension 4, the four-parameter versal deformation  $\mathcal{V}(p) = A - \lambda B + Z_A - \lambda Z_B$ , where  $Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$ , is given by

$$Z_A = \left[ \begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ \hline p_1 & p_1 \bar{\gamma} & p_3(|\gamma|^2 + 1) & 0 \\ p_2 - p_1 \frac{2\gamma}{|\gamma|^2 + 1} & p_2 \bar{\gamma} - p_1 \frac{|\gamma|^2 - 1}{|\gamma|^2 + 1} & -p_3 \gamma + p_4 & p_3(|\gamma|^2 + 1) \end{array} \right]$$

and

$$Z_B = \left[ \begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ \hline -p_1 \bar{\gamma} & -p_1 \bar{\gamma}^2 & -p_3(|\gamma|^2 \bar{\gamma} + \bar{\gamma}) & 0 \\ -p_2 \bar{\gamma} + p_1 \frac{|\gamma|^2 - 1}{|\gamma|^2 + 1} & -p_2 \bar{\gamma}^2 - p_1 \frac{2\bar{\gamma}}{|\gamma|^2 + 1} & -p_3 - p_4 \bar{\gamma} & -p_3(|\gamma|^2 \bar{\gamma} + \bar{\gamma}) \end{array} \right].$$

For the  $11 \times 11$  pencil  $A - \lambda B = L_1 \oplus J_3(0) \oplus N_4 \oplus L_2^T$  with codimension 26, the 26-parameter versal deformation  $\mathcal{V}(p) = A - \lambda B + Z_A - \lambda Z_B$ , where  $Z_A - \lambda Z_B \in$

$\text{nor}(A - \lambda B)$ , is given by

$$Z_A = \left[ \begin{array}{cc|ccc|cccc|cc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_1 & 0 & p_{13} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_2 & p_1 & p_{14} & p_{13} & 0 & 0 & 0 & 0 & 0 & 0 \\ p_3 & p_2 & p_{15} & p_{14} & p_{13} & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & p_4 & 0 & 0 & 0 & p_{19} & 0 & 0 & 0 & 0 \\ p_4 & p_5 & 0 & 0 & 0 & p_{20} & p_{19} & 0 & 0 & 0 \\ p_5 & p_6 & 0 & 0 & 0 & p_{21} & p_{20} & p_{19} & 0 & 0 \\ \hline p_8 & p_9 & p_{18} & p_{17} & p_{16} & p_{23} & 0 & 0 & 0 & 0 \\ p_9 & p_{10} & p_{17} & p_{16} & 0 & p_{24} & p_{23} & 0 & 0 & 0 \\ p_{10} & p_{11} & p_{16} & 0 & 0 & p_{25} & p_{24} & p_{23} & 0 & 0 \end{array} \right]$$

and

$$Z_B = \left[ \begin{array}{cc|ccc|cccc|cc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -p_1 & 0 & -p_{13} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -p_2 & -p_1 & -p_{14} & -p_{13} & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & -p_4 & 0 & 0 & 0 & -p_{19} & 0 & 0 & 0 & 0 \\ -p_4 & -p_5 & 0 & 0 & 0 & -p_{20} & -p_{19} & 0 & 0 & 0 \\ -p_5 & -p_6 & 0 & 0 & 0 & -p_{21} & -p_{20} & -p_{19} & 0 & 0 \\ -p_6 & -p_7 & 0 & 0 & 0 & -p_{22} & -p_{21} & -p_{20} & -p_{19} & 0 \\ \hline -p_9 & -p_{10} & -p_{17} & -p_{16} & 0 & -p_{24} & -p_{23} & 0 & 0 & 0 \\ -p_{10} & -p_{11} & -p_{16} & 0 & 0 & -p_{25} & -p_{24} & -p_{23} & 0 & 0 \\ -p_{11} & -p_{12} & 0 & 0 & 0 & -p_{26} & -p_{25} & -p_{24} & -p_{23} & 0 \end{array} \right].$$

**6.2. Versal deformations of the set of  $2 \times 3$  matrix pencils.** In [15], the algebraic and geometric characteristics of the set of  $2 \times 3$  matrix pencils were examined in full detail, including the complete closure hierarchy. There, all nonzero and finite eigenvalues were considered as unspecified.  $R_2$  was used to denote a  $2 \times 2$  block with nonzero finite eigenvalues, i.e., any of the three structures  $J_1(\alpha) \oplus J_1(\beta)$ ,  $J_1(\alpha) \oplus J_1(\alpha)$ , and  $J_2(\alpha)$ , where  $\alpha, \beta \neq \{0, \infty\}$ . However, in the context of versal deformations all these forms are considered separately and with the eigenvalues specified (known). Consequently, we now have 20 different Kronecker structures to investigate. For example, the versal deformation of  $A - \lambda B = L_0 \oplus J_2(\gamma)$ ,  $\gamma \neq \{0, \infty\}$ , is found by computing  $Z_A - \lambda Z_B =$

(6.1)

$$\left[ \begin{array}{ccc} p_1 + \lambda \bar{\gamma} p_1 & p_3(|\gamma|^2 + 1) + p_3(|\gamma|^2 \bar{\gamma} + \bar{\gamma}) & 0 \\ p_2 - \frac{p_1 \gamma}{|\gamma|^2 + 1} + \lambda(p_2 \bar{\gamma} + \frac{p_1}{|\gamma|^2 + 1}) & -p_3 \gamma + p_4 + \lambda(p_3 + p_4 \bar{\gamma}) & p_3(|\gamma|^2 + 1) + p_3(|\gamma|^2 \bar{\gamma} + \bar{\gamma}) \end{array} \right].$$

In Table 6.1 we show the versal deformations for all different Kronecker structures for this set of matrix pencils. The different structures are displayed in increasing codimension order.

**6.2.1. Using GUPTRI in a random walk in tangent and normal directions of nongeneric pencils.** To illustrate how perturbations in the tangent space and the normal space affect the Kronecker structure computed by a staircase algorithm, we have performed a set of tests on nongeneric  $2 \times 3$  matrix pencils. Since the

TABLE 6.1  
 Versal deformations  $\mathcal{V}(p) = A - \lambda B + Z_A - \lambda Z_B$  of the set of  $2 \times 3$  matrix pencils.

KCF	$A - \lambda B$	$Z_A - \lambda Z_B$
$L_2$	$\begin{bmatrix} -\lambda & 1 & 0 \\ 0 & -\lambda & 1 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$
$L_1 \oplus J_1(\gamma)$	$\begin{bmatrix} -\lambda & 1 & 0 \\ 0 & 0 & \gamma - \lambda \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 0 \\ p_1 + \lambda \bar{\gamma} p_1 & \bar{\gamma} p_1 + \lambda \bar{\gamma}^2 p_1 & p_2 + \lambda \bar{\gamma} p_2 \end{bmatrix}$
$L_1 \oplus J_1(0)$	$\begin{bmatrix} -\lambda & 1 & 0 \\ 0 & 0 & -\lambda \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 0 \\ p_1 & 0 & p_2 \end{bmatrix}$
$L_1 \oplus N_1$	$\begin{bmatrix} -\lambda & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 0 \\ 0 & \lambda p_1 & \lambda p_2 \end{bmatrix}$
$L_0 \oplus J_1(\gamma_1) \oplus J_1(\gamma_2)$	$\begin{bmatrix} 0 & \gamma_1 - \lambda & 0 \\ 0 & 0 & \gamma_2 - \lambda \end{bmatrix}$	$\begin{bmatrix} p_1 + \lambda \bar{\gamma}_1 p_1 & p_3 + \lambda \bar{\gamma}_1 p_3 & 0 \\ p_2 + \lambda \bar{\gamma}_2 p_2 & 0 & p_4 + \lambda \bar{\gamma}_2 p_4 \end{bmatrix}$
$L_0 \oplus J_2(\gamma)$	$\begin{bmatrix} 0 & \gamma - \lambda & 1 \\ 0 & 0 & \gamma - \lambda \end{bmatrix}$	See (6.1)
$L_0 \oplus 2J_1(\gamma)$	$\begin{bmatrix} 0 & \gamma - \lambda & 0 \\ 0 & 0 & \gamma - \lambda \end{bmatrix}$	$\begin{bmatrix} p_1 + \lambda \bar{\gamma} p_1 & p_3 + \lambda \bar{\gamma} p_3 & p_5 + \lambda \bar{\gamma} p_5 \\ p_2 + \lambda \bar{\gamma} p_2 & p_4 + \lambda \bar{\gamma} p_4 & p_6 + \lambda \bar{\gamma} p_6 \end{bmatrix}$
$L_0 \oplus J_1(0) \oplus J_1(\gamma)$	$\begin{bmatrix} 0 & -\lambda & 0 \\ 0 & 0 & \gamma - \lambda \end{bmatrix}$	$\begin{bmatrix} p_1 & p_3 & 0 \\ p_2 + \lambda \bar{\gamma} p_2 & 0 & p_4 + \lambda \bar{\gamma} p_4 \end{bmatrix}$
$L_0 \oplus J_1(\gamma) \oplus N_1$	$\begin{bmatrix} 0 & \gamma - \lambda & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} p_1 + \lambda \bar{\gamma} p_1 & p_3 + \lambda \bar{\gamma} p_3 & 0 \\ \lambda p_2 & 0 & \lambda p_4 \end{bmatrix}$
$L_0 \oplus J_2(0)$	$\begin{bmatrix} 0 & -\lambda & 1 \\ 0 & 0 & -\lambda \end{bmatrix}$	$\begin{bmatrix} p_1 & p_3 & 0 \\ p_2 + \lambda p_1 & p_4 + \lambda p_3 & p_3 \end{bmatrix}$
$L_0 \oplus N_2$	$\begin{bmatrix} 0 & 1 & -\lambda \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} \lambda p_1 & \lambda p_3 & 0 \\ p_1 + \lambda p_2 & p_3 + \lambda p_4 & \lambda p_3 \end{bmatrix}$
$L_0 \oplus J_1(0) \oplus N_1$	$\begin{bmatrix} 0 & -\lambda & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} p_1 & p_3 & 0 \\ \lambda p_2 & 0 & \lambda p_4 \end{bmatrix}$
$L_0 \oplus L_1 \oplus L_0^T$	$\begin{bmatrix} 0 & -\lambda & 1 \\ 0 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & 0 \\ p_1 + \lambda p_2 & p_3 + \lambda p_4 & p_4 + \lambda p_5 \end{bmatrix}$
$L_0 \oplus 2J_1(0)$	$\begin{bmatrix} 0 & -\lambda & 0 \\ 0 & 0 & -\lambda \end{bmatrix}$	$\begin{bmatrix} p_1 & p_3 & p_5 \\ p_2 & p_4 & p_6 \end{bmatrix}$
$L_0 \oplus 2N_1$	$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} \lambda p_1 & \lambda p_3 & \lambda p_5 \\ \lambda p_2 & \lambda p_4 & \lambda p_6 \end{bmatrix}$
$2L_0 \oplus L_1^T$	$\begin{bmatrix} 0 & 0 & -\lambda \\ 0 & 0 & 1 \end{bmatrix}$	$\begin{bmatrix} p_1 + \lambda p_2 & p_4 + \lambda p_5 & 0 \\ p_2 + \lambda p_3 & p_5 + \lambda p_6 & 0 \end{bmatrix}$
$2L_0 \oplus J_1(\gamma) \oplus L_0^T$	$\begin{bmatrix} 0 & 0 & \gamma - \lambda \\ 0 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} p_1 + \lambda \bar{\gamma} p_1 & p_4 + \lambda \bar{\gamma} p_4 & p_7 + \lambda \bar{\gamma} p_7 \\ p_2 + \lambda p_3 & p_5 + \lambda p_6 & p_8 + \lambda \bar{\gamma} p_8 \end{bmatrix}$
$2L_0 \oplus J_1(0) \oplus L_0^T$	$\begin{bmatrix} 0 & 0 & -\lambda \\ 0 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} p_1 & p_4 & p_7 \\ p_2 + \lambda p_3 & p_5 + \lambda p_6 & p_8 \end{bmatrix}$
$2L_0 \oplus N_1 \oplus L_0^T$	$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} \lambda p_1 & \lambda p_4 & \lambda p_7 \\ p_2 + \lambda p_3 & p_5 + \lambda p_6 & \lambda p_8 \end{bmatrix}$
$3L_0 \oplus 2L_0^T$	$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$	$\begin{bmatrix} p_1 + \lambda p_2 & p_5 + \lambda p_6 & p_9 + \lambda p_{10} \\ p_3 + \lambda p_4 & p_7 + \lambda p_8 & p_{11} + \lambda p_{12} \end{bmatrix}$

staircase algorithm considers all nonzero finite eigenvalues as unspecified, we have not included these cases in the test.

For the remaining 12 nongeneric cases a random perturbation  $E_A - \lambda E_B$ , with entries  $e_{ij}^a, e_{ij}^b$ , has been decomposed into two parts  $T_A - \lambda T_B \in \tan(A - \lambda B)$  and

$Z_A - \lambda Z_B \in \text{nor}(A - \lambda B)$  such that

$$E_A = T_A + Z_A \quad \text{and} \quad E_B = T_B + Z_B.$$

We illustrate the decomposition of  $E_A - \lambda E_B$  with  $A - \lambda B = L_0 \oplus J_2(0)$ . From Table 6.1 we get

$$Z_A = \begin{bmatrix} p_1 & p_3 & 0 \\ p_2 & p_4 & p_3 \end{bmatrix}, \quad Z_B = \begin{bmatrix} 0 & 0 & 0 \\ -p_1 & -p_3 & 0 \end{bmatrix}.$$

Let  $T_A - \lambda T_B = (E_A - \lambda E_B) - (Z_A - \lambda Z_B)$ . Now, the parameters  $p_i$  are determined by computing the component of  $E_A - \lambda E_B$  in each of the four orthogonal (but not orthonormal) directions that span the normal space:

$$\begin{aligned} Z_1 &= \frac{1}{2} \left( \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} - \lambda \begin{bmatrix} 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} \right), \\ Z_2 &= 1 \left( \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} - \lambda \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right), \\ Z_3 &= \frac{1}{3} \left( \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix} \right), \\ Z_4 &= 1 \left( \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} - \lambda \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right). \end{aligned}$$

We conclude that

$$p_1 = \frac{e_{11}^a - e_{21}^b}{2}, \quad p_2 = e_{21}^a, \quad p_3 = \frac{e_{12}^a + e_{23}^a - e_{22}^b}{3}, \quad p_4 = e_{22}^a.$$

It is easily verified that  $\langle T_A - \lambda T_B, Z_A - \lambda Z_B \rangle = 0$ .

GUPTRI [11, 12] has been used to compute the Kronecker structure of the perturbed pencils  $A - \lambda B + \epsilon(E_A - \lambda E_B)$ ,  $A - \lambda B + \epsilon(Z_A - \lambda Z_B)$ , and  $A - \lambda B + \epsilon(T_A - \lambda T_B)$  for  $\epsilon = 10^{-16}, 10^{-15}, \dots, 10^0$ . We investigate how far we can move in the tangent and normal directions before GUPTRI reports the generic Kronecker structure.

The procedure has been repeated for all cases and for 100 random perturbations  $(E_A, E_B)$ , where  $\|(E_A, E_B)\|_F = 1$  and  $\|E_A\|_F = \|E_B\|_F$ . The entries of  $(E_A, E_B)$  are uniformly distributed in  $(-0.5, 0.5)$ . For each case and for each perturbation  $E_A - \lambda E_B$  we record the size of  $\epsilon$  when GUPTRI reports the generic Kronecker structure. In Table 6.2 we display the smallest, median, and maximum values of  $\epsilon$  for the 100 random perturbations.

Entries marked + in Table 6.2 mean that the generic structure was not found for any size of the perturbations. All these results were for perturbations in  $\text{tan}(A - \lambda B)$ , and they indicate that for these Kronecker structures there is little or no curvature in the orbit at this point (pencil) in this direction. Here the tangent directions are very close to  $\text{orbit}(A - \lambda B)$ .

Notably, the results for the perturbations  $\epsilon(E_A - \lambda E_B)$  are, except for one case, similar to the results for  $\epsilon(Z_A - \lambda Z_B)$ . This is natural since the perturbation  $E_A - \lambda E_B$  implies a translation both in the tangent space and the normal space directions. The structure changes appear more rapidly in the normal space, i.e., for smaller  $\epsilon$ . Our computational results extend the cone example in section 1.3 to  $2 \times 3$  matrix pencils.

Why is the smallest perturbation  $10^{-16}(Z_A - \lambda Z_B)$  sufficient to find the generic structure for the three cases  $L_0 \oplus 2J_1(0)$ ,  $L_0 \oplus 2N_1$ , and  $3L_0 \oplus 2L_0^T$ ? The explanation is connected to the procedure for determining the numerical rank of matrices.

TABLE 6.2

How far we can move in tangent and normal directions before nongeneric  $2 \times 3$  matrix pencils turn generic.

$A - \lambda B$	$\text{cod}(A - \lambda B)$	$\epsilon(Z_A - \lambda Z_B)$			$\epsilon(T_A - \lambda T_B)$		
		$\epsilon_{\min}$	$\epsilon_{\text{median}}$	$\epsilon_{\max}$	$\epsilon_{\min}$	$\epsilon_{\text{median}}$	$\epsilon_{\max}$
$L_1 \oplus J_1(0)$	2	$10^{-4}$	$10^{-4}$	$10^{-3}$	$10^{-2}$	$10^{-1}$	$10^{-1}$
$L_1 \oplus N_1$	2	$10^{-4}$	$10^{-4}$	$10^{-3}$	$10^{-2}$	$10^{-1}$	$10^0$
$L_0 \oplus J_2(0)$	4	$10^{-4}$	$10^{-4}$	$10^{-3}$	$10^{-2}$	$10^{-1}$	$10^0$
$L_0 \oplus N_2$	4	$10^{-5}$	$10^{-4}$	$10^{-3}$	$10^{-2}$	$10^{-1}$	$10^{-1}$
$L_0 \oplus J_1(0) \oplus N_1$	4	$10^{-4}$	$10^{-4}$	$10^{-2}$	$10^{-2}$	$10^{-1}$	$10^0$
$L_0 \oplus L_1 \oplus L_0^T$	5	$10^{-4}$	$10^{-4}$	$10^{-2}$	$10^{-2}$	$10^{-1}$	$10^0$
$L_0 \oplus 2J_1(0)$	6	$10^{-16}$	$10^{-16}$	$10^{-16}$	+	+	+
$L_0 \oplus 2N_1$	6	$10^{-16}$	$10^{-16}$	$10^{-16}$	+	+	+
$2L_0 \oplus L_1^T$	6	$10^{-4}$	$10^{-4}$	$10^{-2}$	+	+	+
$2L_0 \oplus J_1(0) \oplus L_0^T$	8	$10^{-5}$	$10^{-4}$	$10^{-1}$	+	+	+
$2L_0 \oplus N_1 \oplus L_0^T$	8	$10^{-4}$	$10^{-4}$	$10^{-3}$	+	+	+
$3L_0 \oplus 2L_0^T$	12	$10^{-16}$	$10^{-16}$	$10^{-16}$	+	+	+

GUPTRI has two input parameters, EPSU and GAP, which are used to make rank decisions to determine the Kronecker structure of an input pencil  $A - \lambda B$ . Inside GUPTRI the absolute tolerances  $\text{EPSUA} = \|A\|_E \cdot \text{EPSU}$  and  $\text{EPSUB} = \|B\|_E \cdot \text{EPSU}$  are used in all rank decisions, where the matrices  $A$  and  $B$ , respectively, are involved. Suppose the singular values of  $A$  are computed in increasing order, i.e.,  $0 \leq \sigma_1 \leq \sigma_2 \leq \dots \leq \sigma_k \leq \sigma_{k+1} \leq \dots$ ; then all singular values  $\sigma_k < \text{EPSUA}$  are interpreted as zeros. The rank decision is made more robust in practice: if  $\sigma_k < \text{EPSUA}$  but  $\sigma_{k+1} \geq \text{EPSUA}$ , GUPTRI insists on a gap between the two singular values such that  $\sigma_{k+1}/\sigma_k \geq \text{GAP}$ . If  $\sigma_{k+1}/\sigma_k < \text{GAP}$ ,  $\sigma_{k+1}$  is also treated as zero. This process is repeated until an appreciable gap between the zero and nonzero singular values is obtained. In all of our tests we have used  $\text{EPSU} = 10^{-8}$  and  $\text{GAP} = 1000.0$ .

For the most nongeneric case  $3L_0 \oplus 2L_0^T$ , both the A-part and the B-part are zero matrices giving  $\text{EPSUA} = \text{EPSUB} = 0$ , which in turn leads to the decision that a full rank perturbation  $E_A - \lambda E_B$  times a very small  $\epsilon$  is interpreted as a generic pencil. For the other two cases, either the A-part or the B-part is full rank and the other part is a zero matrix, which accordingly is interpreted to have full rank already for the smallest perturbation.

**6.2.2. Versal deformations and minimal perturbations for changing a nongeneric structure.** In the following we illustrate how versal deformations are useful in the understanding of the relations between the different structures by looking at requirements on perturbations to  $(A, B)$  for changing the Kronecker structure. Assume that we have the following matrix pencil with the Kronecker structure  $L_1 \oplus J_1(0)$ :

$$(6.2) \quad A - \lambda B = \begin{bmatrix} -\epsilon_1 \lambda & \epsilon_2 & 0 \\ 0 & 0 & -\epsilon_3 \lambda \end{bmatrix} \quad \text{and} \quad Z_A - \lambda Z_B = \begin{bmatrix} 0 & 0 & 0 \\ p_1 & 0 & p_2 \end{bmatrix}.$$

It was shown in [15] that  $L_1 \oplus J_1(0)$  with codimension 2 is in the closure of  $\text{orbit}(L_1 \oplus J_1(\gamma))$  ( $\gamma \neq \{0, \infty\}$  but otherwise unspecified) with codimension 1, which in turn is in the closure of  $\text{orbit}(L_2)$  (the generic KCF) with codimension 0. Notice that in Table 6.1, since  $\gamma$  is assumed specified,  $L_1 \oplus J_1(\gamma)$  has two parameters (and codimension = 2). In the discussion that follows we assume that  $\gamma$  is finite and nonzero but unspecified.

We will now, for this example, illustrate how perturbations in the normal space directions can be used to find more generic Kronecker structures (going upward in the Kronecker structure hierarchy) and how we can perturb the elements in  $A - \lambda B$  to find less generic matrix pencils. Since the space spanned by  $Z_A - \lambda Z_B$  is the normal space, we must always first hit a more generic pencil when we move infinitesimally in normal space directions.

The KCF remains unchanged as long as  $p_1 = p_2 = 0$ , but for  $p_1 = 0$  and  $p_2 \neq 0$ , the KCF is changed into  $L_1 \oplus J_1(\gamma)$  (with  $\gamma = p_2$ ). That is, by adding a component in a normal space direction, we find a more generic pencil in the closure hierarchy. Notably, the size of the required perturbation is equal to the smallest size of an eigenvalue to be interpreted as nonzero. By choosing  $p_1$  nonzero (and  $p_2$  arbitrary), the resulting pencil will be generic with the KCF  $L_2$ .

To find a less generic structure, we may proceed in one of the following ways.

1. Find a less generic structure in the closure of  $\text{orbit}(L_1 \oplus J_1(0))$ .
2. Go upward in the closure hierarchy to a more generic structure and then look in that orbit's closure for a less generic structure.

We know from the investigation in [15] that all structures with higher codimension than  $A - \lambda B = L_1 \oplus J_1(0)$  include an  $L_0$  block in their Kronecker structures, which in turn implies that  $A$  and  $B$  must have a common column nullspace of at least dimension 1. Therefore, the smallest perturbation that turns  $L_1 \oplus J_1(0)$  less generic is the smallest perturbation that reduces the rank of

$$\begin{bmatrix} A \\ B \end{bmatrix} = \begin{bmatrix} 0 & \epsilon_2 & 0 \\ 0 & 0 & 0 \\ \epsilon_1 & 0 & 0 \\ 0 & 0 & \epsilon_3 \end{bmatrix}.$$

The size of the smallest rank-reducing perturbation is equal to the smallest of the singular values  $\epsilon_1, \epsilon_2$ , and  $\epsilon_3$ . By just deleting one  $\epsilon_i$ , the corresponding perturbed pencil is a less generic pencil within the closure of  $\text{orbit}(L_1 \oplus J_1(0))$ . These three cases correspond to approach 1 above. We summarize these perturbations and the perturbations in the normal space in Table 6.3. Notice that approach 2 will always require a perturbation larger than  $\min\{\epsilon_i\}$ .

Which of the nongeneric structures displayed in Table 6.3 is obtained by the smallest perturbation to  $L_1 \oplus J_1(0)$ ? Mathematically, it is easy to see that the perturbations in the normal space always can be made smaller than a rank-reducing perturbation  $\epsilon_i$ , since  $p_1$  and  $p_2$  are parameters that can be chosen arbitrarily small, e.g., smaller than  $\min\{\epsilon_i\}$ .

However, in finite-precision arithmetic, it is not clear that the smallest perturbation required to find another structure is in the normal direction. This can be illustrated by using GUPTRI to compute the Kronecker structures for  $A - \lambda B$  as in (6.2) and perturbed as in Table 6.3. For  $\text{EPSU} = 10^{-8}$ ,  $\epsilon_2 = 1$ , and  $\epsilon_1 = \epsilon_3 = 10^{-10}$ , GUPTRI uses different tolerances  $\text{EPSUA} = 10^{-8}$  and  $\text{EPSUB} = 10^{-18}$  for making rank decisions in  $A$  and  $B$ , respectively. It follows that for  $p_1$  and  $p_2$  of order  $10^{-6}$ , GUPTRI still computes the Kronecker structure  $L_1 \oplus J_1(0)$ . However, if  $p_1 = p_2 = 0$  and the B-part of the pencil is perturbed by  $\epsilon_1$  or  $\epsilon_3$ , GUPTRI computes the less generic structures, just as shown in Table 6.3.

**7. Conclusions.** In this paper, we have obtained not only versal deformations for deformations of KCFs, but more importantly for our purposes, metrical information for the perturbation theory of matrix pencils relevant to the KCF. We demon-

TABLE 6.3

*Perturbing  $A - \lambda B$  (defined in (6.2)) yields the pencil  $\tilde{A} - \lambda \tilde{B}$  with more or less generic structures. The codimension of the original orbit is 2.*

$\ (\Delta A, \Delta B)\ _F$	$\tilde{A} - \lambda \tilde{B}$	KCF	$\text{cod}(\tilde{A} - \lambda \tilde{B})$
$p_1$	$\begin{bmatrix} -\epsilon_1 \lambda & \epsilon_2 & 0 \\ p_1 & 0 & -\epsilon_3 \lambda \end{bmatrix}$	$L_2$	0
$p_2$	$\begin{bmatrix} -\epsilon_1 \lambda & \epsilon_2 & 0 \\ 0 & 0 & p_2 - \epsilon_3 \lambda \end{bmatrix}$	$L_1 \oplus J_1(p_2)$	1 (2)
$\epsilon_1$	$\begin{bmatrix} 0 & \epsilon_2 & 0 \\ 0 & 0 & -\epsilon_3 \lambda \end{bmatrix}$	$L_0 \oplus J_1(0) \oplus N_1$	4
$\epsilon_3$	$\begin{bmatrix} -\epsilon_1 \lambda & \epsilon_2 & 0 \\ 0 & 0 & 0 \end{bmatrix}$	$L_0 \oplus L_1 \oplus L_0^T$	5
$\epsilon_2$	$\begin{bmatrix} -\epsilon_1 \lambda & 0 & 0 \\ 0 & 0 & -\epsilon_3 \lambda \end{bmatrix}$	$L_0 \oplus 2J_1(0)$	6

strated with numerical experiments in section 6 how this theory may be used in practice to see how computations are influenced by the geometry. In Part II of this paper, we will explore the stratification theory of matrix pencils with the goal of making algorithmic use of the lattice of orbits under the closure relationship [14].

**Acknowledgments.** We would like to thank Jim Demmel for conveying the message that the geometry of matrix and matrix pencil spaces influence perturbation theory and numerical algorithms. We further thank him, Richard Stanley, and David Vogan for many helpful discussions. In addition, the first author would like to thank the second two authors for their kind hospitality and support at Umeå University where this work and fruitful collaboration began.

REFERENCES

- [1] V. I. ARNOLD, *On matrices depending on parameters*, Russian Math. Surveys, 26 (1971), pp. 29–43.
- [2] T. BEELEN AND P. VAN DOOREN, *An improved algorithm for the computation of Kronecker’s canonical form of a singular pencil*, Linear Algebra Appl., 105 (1988), pp. 9–65.
- [3] J. M. BERG AND H. G. KWATNY, *A canonical parameterization of the Kronecker form of a matrix pencil*, Automatica, 31 (1995), pp. 669–680.
- [4] T. BRÖCKER AND L. LANDER, *Differential Germs and Catastrophes*, Cambridge University Press, London, UK, 1975.
- [5] J. W. BRUCE AND P. J. GIBLIN, *Curves and Singularities*, Cambridge University Press, London, UK, 1991.
- [6] S.-N. CHOW, C. LI, AND D. WANG, *Normal Forms and Bifurcation of Planar Vector Fields*, Cambridge University Press, London, UK, 1994.
- [7] J. DEMMEL, *On structured singular values*, in Proc. of the 27th Conference on Decision and Control, Austin, TX, IEEE, 1988.
- [8] J. DEMMEL AND A. EDELMAN, *The dimension of matrices (matrix pencils) with given Jordan (Kronecker) canonical forms*, Linear Algebra Appl., 230 (1995), pp. 61–87.
- [9] J. DEMMEL AND B. KÅGSTRÖM, *Computing stable eigendecompositions of matrix pencils*, Linear Algebra Appl., 88/89 (1987), pp. 139–186.
- [10] J. DEMMEL AND B. KÅGSTRÖM, *Accurate solutions of ill-posed problems in control theory*, SIAM J. Matrix Anal. Appl., 9 (1988), pp. 126–145.

- [11] J. DEMMEL AND B. KÅGSTRÖM, *The generalized Schur decomposition of an arbitrary pencil  $A - \lambda B$ : Robust software with error bounds and applications. Part I: Theory and algorithms*, ACM Trans. Math. Software, 19 (1993), pp. 160–174.
- [12] J. DEMMEL AND B. KÅGSTRÖM, *The generalized Schur decomposition of an arbitrary pencil  $A - \lambda B$ : Robust software with error bounds and applications. Part II: Software and applications*, ACM Trans. Math. Software, 19 (1993), pp. 175–201.
- [13] A. EDELMAN AND H. MURAKAMI, *Polynomial roots from companion matrix eigenvalues*, Math. Comp., 64 (1995), pp. 763–776.
- [14] A. EDELMAN, E. ELMROTH, AND B. KÅGSTRÖM, *A Geometric Approach to Perturbation Theory of Matrices and Matrix Pencils. Part II: A Stratification-Enhanced Staircase Algorithm*, Report UMINF-96.13, Department of Computing Science, Umeå University, Umeå, Sweden, 1996; SIAM J. Matrix Anal. Appl., submitted.
- [15] E. ELMROTH AND B. KÅGSTRÖM, *The set of 2-by-3 matrix pencils—Kronecker structures and their transitions under perturbations*, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 1–34.
- [16] F. GANTMACHER, *The Theory of Matrices, Vol. I and II*, translation, Chelsea, NY, 1959.
- [17] G. GOLUB AND C. VAN LOAN, *Matrix Computations*, 2nd ed., The Johns Hopkins University Press, Baltimore, MD, 1989.
- [18] M. GOLUBITSKY AND V. GUILLEMIN, *Stable Mappings and their Singularities*, Springer-Verlag, Berlin, New York, 1973.
- [19] U. HELMKE AND D. HINRICHSSEN, *Canonical forms and orbit spaces of linear systems*, IMA J. Math. Control Inform., 3 (1986), pp. 167–184.
- [20] D. HINRICHSSEN AND D. PRÄTZEL-WOLTERS, *A wild quiver in linear systems theory*, Linear Algebra Appl., 91 (1987), pp. 143–175.
- [21] D. HINRICHSSEN AND D. PRÄTZEL-WOLTERS, *A Jordan canonical form for reachable linear systems*, Linear Algebra Appl., 122/123/124 (1989), pp. 489–524.
- [22] B. KÅGSTRÖM, *RGSVD—an algorithm for computing the Kronecker canonical form and reducing subspaces of singular matrix pencils  $A - \lambda B$* , SIAM J. Sci. Stat. Comp., 7 (1986), pp. 185–211.
- [23] V. B. KHAZANOV AND V. KUBLANOVSKAYA, *Spectral problems for matrix pencils. Methods and algorithms. I*, Soviet J. Numer. Anal. Math. Modelling, 3 (1988), pp. 337–371.
- [24] V. KUBLANOVSKAYA, *AB-algorithm and its modifications for the spectral problem of linear pencils of matrices*, Numer. Math., 43 (1984), pp. 329–342.
- [25] Y.-C. LU, *Singularity Theory and an Introduction to Catastrophe Theory*, Springer-Verlag, Berlin, New York, 1976.
- [26] A. POKRZYWA, *On perturbations and the equivalence orbit of a matrix pencil*, Linear Algebra Appl., 82 (1986), pp. 99–121.
- [27] D. PRÄTZEL-WOLTERS, *Canonical forms for linear systems*, Linear Algebra Appl., 50 (1983), pp. 437–473.
- [28] M. SHUB AND S. SMALE, *Complexity of Bezout’s theorem II: Volumes and probabilities*, in Computational Algebraic Geometry, F. Eyssette and A. Galligo, eds., Progress in Mathematics 109, Birkhauser, 1993, pp. 267–285.
- [29] G. W. STEWART AND J.-G. SUN, *Matrix Perturbation Theory*, Academic Press, New York, 1990.
- [30] L. STOLOVITCH, *On the computation of a versal family of matrices*, Numer. Algorithms, 4 (1993), pp. 25–56.
- [31] P. VAN DOOREN, *The computation of Kronecker’s canonical form of a singular pencil*, Linear Algebra Appl., 27 (1979), pp. 103–141.
- [32] P. VAN DOOREN, *The generalized eigenstructure problem in linear system theory*, IEEE Trans. Automat. Control, AC-26 (1981), pp. 111–129.
- [33] P. VAN DOOREN, *Reducing subspaces: Definitions, properties and algorithms*, in Matrix Pencils, B. Kågström and A. Ruhe, eds., Springer-Verlag, Berlin, 1983, pp. 58–73; Lecture Notes in Mathematics 973, Proceedings, Pite Havsbad, 1982.
- [34] W. WATERHOUSE, *The codimension of singular matrix pairs*, Linear Algebra Appl., 57 (1984), pp. 227–245.
- [35] J. H. WILKINSON, *The Algebraic Eigenvalue Problem*, Oxford Science Publications, Oxford, UK, 1965.
- [36] J. H. WILKINSON, *Linear differential equations and Kronecker’s canonical form*, in Recent Advances in Numerical Analysis, C. de Boor and G. Golub, eds., Academic Press, New York, 1978, pp. 231–265.



Copyright of SIAM Journal on Matrix Analysis & Applications is the property of Society for Industrial and Applied Mathematics and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.